



First integrals for nonlinear dispersive equations

Frédéric Hélein

► **To cite this version:**

| Frédéric Hélein. First integrals for nonlinear dispersive equations. 2013. hal-00879582

HAL Id: hal-00879582

<https://hal-univ-diderot.archives-ouvertes.fr/hal-00879582>

Submitted on 4 Nov 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

First integrals for nonlinear dispersive equations

Frédéric HÉLEIN*

November 4, 2013

Abstract

Given a solution of a semilinear dispersive partial differential equation with a real analytic nonlinearity, we relate its Cauchy data at two different times by nonlinear representation formulas in terms of convergent series. These series are constructed by means of generating functions. All this theory is based on a new suitable formulation of the dynamics of solutions of dispersive equations.

Consider a partial differential equation

$$Lu + N(u, \partial u) = 0, \tag{1}$$

which describes the evolution of a map u from a space-time \mathbb{R}^{n+1} (with coordinates (x^0, \dots, x^n)) to a finite dimensional vector space. Here L is a linear differential operator (e.g. the wave operator $\square = \partial_0^2 - \Delta$, Klein–Gordon $\square + m^2$, Dirac $\not{\partial} + im$, or any combination) and N is a real analytic nonlinear function on u and its first space-time derivatives ∂u . For any $t \in \mathbb{R}$, denote by $[u]_t$ the Cauchy data of u at time t . We address the question: assume that we know $[u]_{t_1}$ for some $t_1 \in \mathbb{R}$, can we compute the value of u at a point at another time t_2 ? If N is a linear function the answer is positive and is given by a linear integral formula, if N is a polynomial this may also work by using series, i.e. an infinite sum of multilinear integrals, as we will present here. In [19] D. Harrivel obtained such a result for a (roughly speaking \mathcal{C}^2) solution of the scalar Klein–Gordon equation $\square u + m^2 u + \lambda u^2 = 0$. It amounts to build a time dependant family of functionals $(\mathcal{S}_t)_t$ of Cauchy data s.t., if u solves (1), then $\mathcal{S}_t([u]_t)$ does not depend on t . Moreover one can prescribe \mathcal{S}_{t_2} to be any linear functional. By choosing e.g. \mathcal{S}_{t_2} to be the Dirac distribution at some point we thus get a positive answer of the previous question. The functionals \mathcal{S}_t are series, each term of which is a sum of integrals over Cartesian products of the space-time built from planar binary trees by using Feynman rules. The important point is that one can ensure that the series converges for $|t_2 - t_1|$ sufficiently small.

*Institut de Mathématiques de Jussieu—Paris Rive Gauche, UMR CNRS 7586, Université Paris Diderot — Paris 7, Bâtiment Sophie Germain, Case 7012, 75205 Paris Cedex 13, France, helein@math.univ-paris-diderot.fr

In [21] this result was further extended to systems with more general (real analytic) nonlinearities and for less regular solutions (roughly speaking \mathcal{C}^1). The method, which was different from [19], did not use a combinatorial analysis of the series, but rests on the construction of a generating function which, by using Wick's theorem for developing it, gives us the desired expansion.

The following paper presents an improvement of the results in [21]. A new ingredient is a different formulation of the dynamics, which allows us to deal with even less regular solutions (roughly speaking \mathcal{C}^0 in general). This formulation is, we believe, new although it is a straightforward consequence of the well-known Duhamel formula. To explain it, consider the standard way to formulate an evolution PDE such as (1) as an ODE in the infinite dimensional space of all Cauchy data:

$$\frac{d[u]_t}{dt} = X([u]_t). \quad (2)$$

We introduce an alternative formulation: we work in the space \mathcal{E}_0 of solutions to Equation (3) below and replace $[u]_t$ by the unique solution φ to the linear equation

$$L\varphi = 0 \quad (3)$$

the *Cauchy data of which is* $[u]_t$, i.e. the same as u at time t . We denote by $\Theta_t u \in \mathcal{E}_0$ this solution. Let G be the homogeneous solution to $LG = 0$ s.t., if $Y : M \rightarrow \mathbb{R}$ is the function defined by $Y(x) = 1$ if $x^0 \geq 0$ and $G(x) = 0$ if $x^0 < 0$, for some time coordinate x^0 , then YG is the retarded fundamental solution of L . Consider the time dependant vector field $(V_t)_t$ on \mathcal{E}_0 defined by $V_t \varphi := \int_{y^0=t} d\vec{y} G(\cdot - y) N(\varphi, \partial\varphi)(y)$. Our first result is:

Theorem 0.1 *A map u is a solution of (1) if and only if the map $t \mapsto \Theta_t u$ is a solution to*

$$\frac{d\Theta_t u}{dt} + V_t(\Theta_t u) = 0. \quad (4)$$

A precise statement of this result is the content of Theorem 2.1. An advantage of Equation (4) is that it is manifestly covariant: the space \mathcal{E}_0 in which $\Theta_t u$ takes values does not depend on t nor on any choice of space-time coordinates, in contrast with the target space of $t \mapsto [u]_t$. This advantage is even more striking on a curved space-time, where a similar result will be proved (Theorem 3.2). A second advantage is that the map $t \mapsto \Theta_t u$ is more regular than $t \mapsto [u]_t$: under general hypotheses, if u is a weak solution of (1) then $t \mapsto \Theta_t u$ is \mathcal{C}^1 !

This formulation is also useful for the problem expounded previously. Consider the space \mathbb{F} of real analytic functionals on \mathcal{E}_0 . We define for all t the first order 'differential' linear operator $V_t \cdot$ acting on \mathbb{F} by:

$$\forall f \in \mathbb{F}, \forall \varphi \in \mathcal{E}_0, \quad (V_t \cdot f)(\varphi) := \delta f_\varphi(V_t(\varphi)), \quad (5)$$

where δf_φ is the differential of f at φ . Then one of our main result is that we can make sense of the chronological exponential $T \exp \left(\int_{t_1}^{t_2} ds V_s \cdot \right)$ as a linear operator acting on \mathbb{F} ,

continuous in a suitable topology, if $|t_2 - t_1|$ is sufficiently small. This operator is the key for constructing the family $(\mathcal{S}_t)_t$ of operators such that $\mathcal{S}_t([u]_t)$ does not depend on t if u is a solution of (1):

Theorem 0.2 *Let $r > 0$. There exists a constant $\bar{t} > 0$ which depends on Equation (1) and on r , such that, for any $t_1, t_2 \in \mathbb{R}$ such that $|t_2 - t_1| < \bar{t}$ and for any $f \in \mathbb{F}$, with a radius of convergence r , the functional*

$$U_{t_1}^{t_2} f := \text{Exp} \left(\int_{t_1}^{t_2} ds V_s \cdot \right) f$$

is well defined on a ball in \mathbb{F} and has a non vanishing radius of convergence R . Moreover, if u is a solution of (1) the Cauchy data of which is smaller than R , then

$$(U_{t_1}^{t_2} f)(\Theta_{t_2} u) \quad \text{is equal to} \quad f(\Theta_{t_1} u). \quad (6)$$

Details on the statement in Theorem 0.2 (the topology on \mathcal{E}_0 and on the space of Cauchy data) will given in the next Section. In general we will set $u \in \mathcal{C}^0(I, H^s(\mathbb{R}^n)) \cap \mathcal{C}^1(I, H^{s-r}(\mathbb{R}^n))$, where r depend on L (e.g. $r = 1$ for $L = \square$) and $s > n/2$ in general. However for a Klein–Gordon equation with some polynomial nonlinearity, it may work for some special values of s and n s.t. $s \leq n/2$ (see Remark 2.1).

This result can be restated in a different language inspired by perturbative quantum fields theory: $U_{t_1}^{t_2} f$ can be written

$$U_{t_1}^{t_2} f = \left(\text{Exp} \int_{t_1 < y^0 < t_2} dy N^i(\phi, \partial\phi)(y) \phi_i^+(y) \right) f,$$

where ϕ and ϕ^+ are kind of *creation* and *annihilation* operators respectively (see Section 7 for details).

Plan of the paper

For simplicity most results are presented for a differential operator with constant coefficients on a flat space-time. Section 1 contains the notations and a precise formulation of the hypotheses needed for the theory on a flat space-time. In Section 2 we construct the map $u \mapsto \Theta_t u$ and the vector field V_t on a flat space-time. We end with the proof of Theorem 2.1, a version of Theorem 0.1 on a flat space-time. We also show that V_t is real analytic on an open ball in \mathcal{E}_0 . In Section 3 we extend these results to a curved space-time. For simplicity we restrict ourself to the Klein–Gordon operator and a cubic nonlinearity. We show also that the formulation (4) works if we replace a foliation by space-like hypersurfaces which are the level sets of a time function by a more general family of space-like hypersurfaces which may overlap.

In Sections 4, 5 and 6 we develop a theory valid in any Banach space \mathbb{X} . In Section 4 we introduce various topologies on the space $\mathbb{F}(\mathbb{X})$ of real analytic functions on bounded balls of \mathbb{X} . We define in particular, for any $r \in [0, +\infty]$, the space $\mathbb{F}_r(\mathbb{X})$ of real analytic functions on \mathbb{X} which, roughly speaking, have a radius of convergence greater or equal to

r. We also derive properties satisfied by a time dependant family $(V_t)_{t \in \mathbb{R}}$ of real analytic first order differential operators acting on $\mathbb{F}(\mathbb{X})$. In Section 5 we prove the existence of the chronological exponential $U_{t_1}^{t_2} = T \exp \left(\int_{t_1}^{t_2} ds V_s \cdot \right)$ as a bounded operator acting between subspaces of $\mathbb{F}(\mathbb{X})$, if $|t_2 - t_1|$ is small enough. The difficulty is that the operators $V_t \cdot$ are not bounded in any topology. Hence the chronological exponential cannot make sense as a bounded operator from a topological to itself. However we will prove that $U_{t_1}^{t_2}$ maps continuously $\mathbb{F}_R(\mathbb{X})$ to $\mathbb{F}_{e^{-|t_2-t_1|X(R)}}(\mathbb{X})$, where X is a (positive) real analytic vector field on \mathbb{R} which is constructed out of Equation (1) and of the choice of topology on the set of its solutions. In Section 6 we prove that $(U_{t_1}^t f)(\varphi(t))$ does not depend on t if φ is a solution of $\frac{d\varphi}{dt} + V_t(\varphi) = 0$, a result which, combined with Theorem 2.1 or Theorem 3.2, implies different versions of Theorem 0.2.

In Section 7 and 8 we give some applications of our results and discuss the analogy and the difference with methods from Quantum Field Theory.

Further comments

This work is motivated by questions in [24, 25]. Our formulation of the dynamics by (24) can be viewed as an analogue for dispersive partial differential equations of Lagrange's method of *variation of the constant*, it is also an infinitesimal version of Duhamel's formula (45). This is the reason for the name 'Lagrange–Duhamel' for V_t .

Developping (6) by using Wick theorem leads to an expansion in terms of 'Feynman trees', as for instance (99). A heuristic way to understand where this comes from consists in inserting the l.h.s. of (47) in the integral in the r.h.s. of it and in iterating this process. Then we see easily that $u(x)$ should be expressed as the sum of a formal series. But it seems difficult to prove directly by this method that this process converges and to estimate the radius of convergence of the series. On the other hand this process is also the key of the Picard fixed point Theorem which is used to prove the local existence of solutions. However the proof of the fixed point Theorem is based on precise estimates of the previous process but it hides the structure of the series which is generated by this process. Our result can hence be understood as filling the gap between both ways.

Series expansions of solution to nonlinear *ordinary* differential equations (ODE) have a long history. We can mention Lie series defined by K.T. Chen [12], the Chen–Fliess series [16] introduced in the framework of control theory by M. Fliess (or some variants like Volterra series or Magnus expansion [29]) which are extensively used in control theory [1, 32, 26] but also in the study of dynamical systems and in numerical analysis. Other major tools are Butcher series which explain the structure of Runge–Kutta methods of approximation of the solution of an ODE. They have been introduced by J.C. Butcher [10] and developed by E. Hairer and G. Wanner [18] which explain that Runge–Kutta methods are governed by trees. Later on C. Brouder [8, 9] realized that the structure which underlies the original Butcher's computation coincides with the Hopf algebra defined by D. Kreimer in his work about the renormalization theory [27]. Concerning analogous results on nonlinear *partial* differential equations, it seems that the fact that one can represent solutions or functionals on the set of solutions by series indexed by trees

is known to physicists since the work of J. Schwinger and R. Feynman (and Butcher was also aware of that in his original work). However it is not that easy to find precise references in the literature: the Reader may look e.g. at [14], where a formal series expansion is presented and the recent paper [15] for comparison with quantum field theory. But, to our knowledge, the only rigorous results (i.e. with a proof of convergence of the series) can be found in [19, 21].

We have used relatively elementary tools from the analysis of PDE's and, in particular, we do not rely on the modern theory for wave and Schrödinger equations (Strichartz estimates, Klainerman bilinear estimates, etc.). Many improvements in these directions could be provided, although they may not be straightforward. Also we are not able to apply our theory the KdV equation, since its nonlinearity cannot be controlled by our methods. Another question concerns the extension of our results to an infinite time interval and to relate together the asymptotic data for $t \rightarrow -\infty$ and $t \rightarrow +\infty$. One may indeed ask whether the limits $u_{\pm} := \lim_{t \rightarrow \pm\infty} \Theta_t u$ exist and, if so, if for $f \in \mathbb{F}$,

$$f(u_-) = \left(\left(T \exp \int_{-\infty}^0 d\tau V_{\tau} \cdot \right) f \right) (\Theta_0 u) = \left(\left(T \exp \int_{-\infty}^{+\infty} d\tau V_{\tau} \cdot \right) f \right) (u_+).$$

Such identities (and their analogues by exchanging u_+ and u_-) would imply in particular that the scattering map $S : u_- \mapsto u_+$ and the wave operators $W_{\pm} : u_{\pm} \mapsto \Theta_0 u$ are well-defined and real analytic¹. In the light of results in [30, 33, 34, 31, 7, 4, 5] this should be true for the equation $\square u + u^3 = 0$ on \mathbb{R}^{1+3} and for $s = 1$, due to dispersive effects (Strichartz estimates). The key point in all these works is an estimate of the type $\int_{-\infty}^{+\infty} \left(\int_{\mathbb{R}^3} u^6 d\vec{x} \right)^{1/2} dt < +\infty$, which, e.g., holds for a solution u of $\square u + u^3 = 0$ with finite energy.

Acknowledgements — I wish to thank Isabelle Gallagher for explanations about paraproducts. This paper is an extended and improved version of an earlier work in collaboration with Dikanaina Harrivel [21].

1 Notations and hypotheses

Generalities — $M := \mathbb{R} \times \mathbb{R}^n$ represents an $(n + 1)$ -dimensional flat space-time. We denote by $x = (x^0, \vec{x}) = (x^0, x^1, \dots, x^n)$ the coordinates on M and set $\partial_{\mu} = \frac{\partial}{\partial x^{\mu}}$ for $0 \leq \mu \leq n$. We let E be a finite dimensional real vector space and we consider maps from M to E .

For any smooth fastly decreasing functions $f \in \mathcal{S}(\mathbb{R}^n)$ we define its Fourier transform $\hat{f}(\xi) = \int_{\mathbb{R}^n} f(\vec{x}) e^{-i\vec{x} \cdot \xi} d\vec{x}$ and we extend it to space $\mathcal{S}'(\mathbb{R}^n)$ of tempered distributions by the standard duality argument. In case of a map f which depends on $(t, \vec{x}) \in I \times \mathbb{R}^n$, we also denote by $\hat{f}(t, \xi) = \int_{\mathbb{R}^n} f(t, \vec{x}) e^{-i\vec{x} \cdot \xi} d\vec{x}$ the partial Fourier transform in space variables.

¹In our definition S and W_{\pm} map \mathcal{E}_0^s to itself. This differs from most references where the scattering map reads in our notations $\Phi_0^{-1} \circ S \circ \Phi_0 : \text{Cau}^s \rightarrow \text{Cau}^s$ and the wave maps are $\Phi_0^{-1} \circ W_{\pm} \circ \Phi_0 : \text{Cau}^s \rightarrow \text{Cau}^s$ (see Paragraph 2.1 for the definition of Φ_0).

For $s \in \mathbb{R}$, we let $H^s(\mathbb{R}^n) := \{\varphi \in \mathcal{S}'(\mathbb{R}^n) \mid [\xi \mapsto \langle \xi \rangle^s \widehat{\varphi}(\xi)] \in L^2(\mathbb{R}^n)\}$, where $\langle \xi \rangle := \sqrt{m^2 + |\xi|^2}$ and we set $\|\varphi\|_{H^s} := \|\langle \xi \rangle^s \widehat{\varphi}\|_{L^2}$. We let $H^s(\mathbb{R}^n, E)$ be the Sobolev space of E -valued maps on \mathbb{R}^n . If $\varphi \in H^s(\mathbb{R}^n, E)$ has the coordinates φ^i ($1 \leq i \leq \dim E$) in a basis of E we set

$$\|\varphi\|_{H^s} := \sum_{i=1}^{\dim E} \|\varphi^i\|_{H^s}. \quad (7)$$

The class of differential operators L — We suppose that there is a splitting $E := E_1 \oplus E_2$, where E_1 and E_2 are two vector subspaces of E . This leads to a decomposition of any map $\varphi : M \rightarrow E$ as $\varphi = (\varphi_1, \varphi_2)$. We assume that the linear differential operator L acting on smooth maps $u : M \rightarrow E$ has the form

$$L = \begin{pmatrix} L_1 & 0 \\ 0 & L_2 \end{pmatrix} = \begin{pmatrix} \gamma^0 \partial_0 + P_1(\vec{\partial}) & 0 \\ 0 & \partial_0^2 + P_2(\vec{\partial}) \end{pmatrix}, \quad (8)$$

where $\gamma^0 \in \text{End}(E_1)$ is an invertible matrix, $\vec{\partial} := (\partial_1, \dots, \partial_n)$ and P_1 and P_2 are polynomials with coefficients in respectively $\text{End}(E_1)$ and $\text{End}(E_2)$ and of degree respectively r and $2r$, where $r \in \mathbb{N}^*$.

We assume that, $\forall \xi \in (\mathbb{R}^n)^*$, $i(\gamma^0)^{-1}P_1(i\xi)$ is a Hermitian matrix and $P_2(i\xi)$ is positive Hermitian. Moreover we suppose that there exists two constants $\alpha > 0$ and $\mu_0 \geq 0$ s.t., in the sense of Hermitian operators acting on E_2 ,

$$\forall \xi \in (\mathbb{R}^n)^*, \quad P_2(i\xi) \geq \alpha(\mu_0 + |\xi|^{2r}). \quad (9)$$

Below is a list of examples for L (setting $\square = \partial_0^2 - \Delta$).

	L	E_1	E_2	r
Klein–Gordon	$\square + m^2$	$\{0\}$	\mathbb{R}	1
Schrödinger	$i\partial_0 + \Delta$	\mathbb{C}	$\{0\}$	2
Dirac on \mathbb{R}^4	$\not{\partial} + im = \gamma^\mu \partial_\mu + im$	\mathbb{C}^4	$\{0\}$	1
linearized Korteweg–de Vries	$\partial_0 + (\partial_1)^3$	\mathbb{R}	$\{0\}$	3
linearized Dirac–Maxwell (in Lorentz gauge)	$\begin{pmatrix} \not{\partial} + im & 0 \\ 0 & \square \end{pmatrix}$	\mathbb{C}^4	\mathbb{R}^4	1

The function spaces — For any $s \in \mathbb{R}$ and any interval $I \subset \mathbb{R}$ we define the space

$$\mathcal{F}^s(I) := \mathcal{C}^0(I, H^s(\mathbb{R}^n, E)) \cap \mathcal{C}^1(I, H^{s-r}(\mathbb{R}^n, E))$$

on which the operator L acts. The natural space of Cauchy data for L on $\mathcal{F}^s(I)$ is $\text{Cau}^s := H^s(\mathbb{R}^n, E) \times H^{s-r}(\mathbb{R}^n, E_2)$. For any $(\psi, \chi) \in \text{Cau}^s$, we set

$$\|\psi, \chi\|_{\text{Cau}^s} := \|\psi\|_{H^s} + \|\chi\|_{H^{s-r}}.$$

The space $\mathcal{F}^s(I)$ is equipped with the norm $\|u\|_{\mathcal{F}^s(I)} := \sup_{\tau \in I} \|[u]_\tau\|_{\text{Cau}^s} = \|u\|_{L^\infty(I, H^s)} + \|\partial_0 u_2\|_{L^\infty(I, H^{s-r})}$.

For any map φ defined on a neighbourhood of $\{t\} \times \mathbb{R}^n$ in M , define its Cauchy data at time t by $[\varphi]_t := (\varphi|_t, \partial_0 \varphi_2|_t) \in \text{Cau}^s$, where, for any function ψ , we note $\psi|_t := \psi(t, \cdot)$ its restriction to $\{t\} \times \mathbb{R}^n$ (which we identify with a function defined on \mathbb{R}^n). For any $I \subset \mathbb{R}$ and $t \in I$, this hence defines a continuous linear map of norm one

$$\begin{aligned} \mathcal{F}^s(I) &\longrightarrow \text{Cau}^s \\ \varphi &\longmapsto [\varphi]_t \end{aligned} \quad (10)$$

For any interval $I \subset \mathbb{R}$ we define the space of solutions to the linear equation $L\varphi = 0$:

$$\mathcal{E}_0^s(I) := \{\varphi \in \mathcal{F}^s(I) \mid L\varphi = 0 \text{ weakly}\}. \quad (11)$$

This space is equipped with the norm $\|u\|_{\mathcal{F}^s(I)}$.

By Proposition 2.2, assuming Hypotheses (8) and (9), for any $t \in \mathbb{R}$ and any pair $(\psi, \chi) \in \text{Cau}^s$, there exists a unique map $\varphi \in \mathcal{E}_0^s(\mathbb{R})$ s.t. $[\varphi]_t = (\psi, \chi)$, i.e. a solution $\varphi \in \mathcal{F}^s(\mathbb{R})$ of:

$$L\varphi = 0 \quad \text{s.t.} \quad \varphi|_t = \psi \quad \text{and} \quad \partial_0 \varphi|_t = \chi. \quad (12)$$

We denote by $\Phi_t(\psi, \chi)$ this solution.

The map Θ — For any map u defined on a neighbourhood of $\{t\} \times \mathbb{R}^n$ we set

$$\Theta_t u := \Phi_t([u]_t),$$

i.e. $\Theta_t u$ is the unique solution φ of $L\varphi = 0$ s.t. $[\varphi]_t = [u]_t$. This hence defines the map

$$\begin{aligned} \Theta : I \times \mathcal{F}^s(I) &\longrightarrow I \times \text{Cau}^s \longrightarrow \mathcal{E}_0^s(\mathbb{R}) \\ (t, u) &\longmapsto (t, [u]_t) \longmapsto \Theta_t u \end{aligned} \quad (13)$$

Polynomials and real analytic functions — Let \mathbb{X}, \mathbb{Y} be two Banach spaces and $p \in \mathbb{N}$. For any $r > 0$, denote by $B_{\mathbb{X}}(r)$ the open ball of radius r and of center 0 in \mathbb{X} . A linear map f_\otimes from $\mathbb{X}^{\otimes p}$ to \mathbb{Y} is *symmetric* if $\forall \varphi_1, \dots, \varphi_p \in \mathbb{X}, \forall \sigma \in \mathfrak{S}(p), f_\otimes(\varphi_{\sigma(1)} \otimes \dots \otimes \varphi_{\sigma(p)}) = f_\otimes(\varphi_1 \otimes \dots \otimes \varphi_p)$, where $\mathfrak{S}(p)$ is the symmetric group with p elements. A *homogeneous polynomial map* $f : \mathbb{X} \rightarrow \mathbb{Y}$ of *degree* p is a map such that there exists a symmetric linear map $f_\otimes : \mathbb{X}^{\otimes p} \rightarrow \mathbb{Y}$ such that $\forall \varphi \in \mathbb{X}, f(\varphi) = f_\otimes(\underbrace{\varphi \otimes \dots \otimes \varphi}_p)$.

Note that f_\otimes , if it exists, is unique and is given by the polarization formula:

$$f_\otimes(\varphi_1 \otimes \dots \otimes \varphi_p) = \frac{1}{2^p p!} \sum_{\epsilon \in \{\pm 1\}^p} \left(\prod_{j=1}^p \epsilon_j \right) f \left(\sum_{j=1}^p \epsilon_j \varphi_j \right). \quad (14)$$

If so we denote by $\|f\|_\otimes$ the smallest nonnegative constant such that $\forall \varphi_1, \dots, \varphi_p \in \mathbb{X}$,

$$\|f_\otimes(\varphi_1 \otimes \dots \otimes \varphi_p)\|_{\mathbb{Y}} \leq \|f\|_\otimes \|\varphi_1\|_{\mathbb{X}} \cdots \|\varphi_p\|_{\mathbb{X}} \quad (15)$$

Most of the time we will abuse notations identifying f_{\otimes} with f , when there is no ambiguity. We denote by $\mathcal{Q}^p(\mathbb{X}, \mathbb{Y})$ the vector space of homogeneous polynomial maps from \mathbb{X} to \mathbb{Y} of degree p .

A *formal series* f from \mathbb{X} to \mathbb{Y} is an infinite sum

$$f = \sum_{p=0}^{\infty} f^{(p)}, \quad (16)$$

where $\forall p \in \mathbb{N}$, $f^{(p)} \in \mathcal{Q}^p(\mathbb{X}, \mathbb{Y})$. The *multiradius of convergence*² of f is the radius of convergence of the series

$$[f](z) := \sum_{p=0}^{\infty} \|f^{(p)}\|_{\otimes} z^p \quad (17)$$

and is denoted by $\rho_{\otimes}(f)$. We denote by $\mathbb{F}(\mathbb{X}, \mathbb{Y})$ the space of formal series from \mathbb{X} to \mathbb{Y} . If $\rho_{\otimes}(f) > 0$, f defines a *real analytic map* from $B_{\mathbb{X}}(\rho_{\otimes}(f))$ to \mathbb{Y} by the relation $\forall \varphi \in B_{\mathbb{X}}(\rho_{\otimes}(f))$, $f(\varphi) = \sum_{p=0}^{\infty} f^{(p)}(\varphi)$. This map is continuous (Lemma 2.2) and satisfies the inequality

$$\|f(\varphi)\|_{\mathbb{Y}} \leq [f](\|\varphi\|_{\mathbb{X}}). \quad (18)$$

For any $r \in (0, +\infty)$, we let $\mathbb{F}_r(\mathbb{X}, \mathbb{Y})$ be the space of formal series f s.t. $[f](r) < +\infty$. In the case where $\mathbb{Y} = \mathbb{R}$, we simply note $\mathbb{F}_r(\mathbb{X}) := \mathbb{F}_r(\mathbb{X}, \mathbb{R})$

Lastly a family $(f_a)_{a \in A}$ of elements in $\mathbb{F}(\mathbb{X}, \mathbb{Y})$ is called a *normal family of analytic maps of multiradius r* if there exists $X \in \mathbb{F}(\mathbb{R})$ s.t. $\rho(X) = r$ and, setting $X(z) = \sum_{p=0}^{\infty} X^{(p)} z^p$, $\forall a \in A$, $\forall p \in \mathbb{N}$, $\|f_a^{(p)}\|_{\otimes} \leq X^{(p)}$ (hence in particular $\rho_{\otimes}(f_a) \geq r$).

The nonlinearity — We note $E_{(1)} := E \times E_2 \times \mathcal{L}(\mathbb{R}^n, E)$. We assume that the map N is real analytic from $E_{(1)}$ to E and that its multiradius of convergence is positive.

For applications to equations in Physics, we are particularly interested in systems (1) of the form³:

$$\begin{cases} L_1 u_1 + N_1(u) & = 0 \\ L_2 u_2 + N_2(u, \partial_0 u_2, \vec{\partial} u) & = 0 \end{cases} \quad (19)$$

where $N_1 : E \rightarrow E_1$ and $N_2 : E_{(1)} \rightarrow E_2$. Motivated by the Yang–Mills system, we are led to consider the case where N_2 is *affine in ∂u* , i.e. there exist real analytic functions J and $K_i^{\mu} : E \rightarrow E_2$ s.t.

$$N_2(u, \partial u) = J(u) + \sum_{i=1}^{\dim E} \sum_{\mu=0}^n K_i^{\mu}(u) \partial_{\mu} u^i. \quad (20)$$

²Note that beside $\|f^{(p)}\|_{\otimes}$ defined by (15), one can also define $\|f^{(p)}\| := \inf_{\varphi \in \mathbb{X} \setminus \{0\}} \|f^{(p)}(\varphi)\|_{\mathbb{Y}} / \|\varphi\|_{\mathbb{X}}^p$ and the radius of convergence $\rho(f)$ of the series $\sum_{p=0}^{\infty} \|f^{(p)}\| z^p$. One can then prove by using (14) that $\|f^{(p)}\| \leq \|f^{(p)}\|_{\otimes} \leq \frac{p!}{p!} \|f^{(p)}\|$, which implies by using Stirling's formula that $e^{-1} \rho(f) \leq \rho_{\otimes}(f) \leq \rho(f)$.

³Actually any system of the form $L_1 u_1 + N_1(u) = 0$ and $L_2 u_2 + \hat{N}_2(u, \partial u) = 0$ can be set in the form (19) through the substitution $N_2(u, \partial_0 u_2, \vec{\partial} u) := \hat{N}_2(u, -N_1(u), \partial_0 u_2, \vec{\partial} u)$.

By setting $N := (N_1, N_2)$, we see that System (19) is equivalent to (1).

For any interval $I \subset \mathbb{R}$, we define

$$\mathcal{E}^s(I) := \{u \in \mathcal{F}^s(I) \mid Lu + N(u, \partial u) = 0 \text{ weakly}\}. \quad (21)$$

The Lagrange–Duhamel vector field — First define the ‘Green function’ G to be the unique distribution on M with coefficients in $\text{End}(E)$, which is a solution of $LG = 0$ and $L(YG) = \delta_0^{1+n} \otimes 1_E$, where $Y(x) := 1_{[0,+\infty)}(x^0)$ is the Heaviside function. Note that through the splitting $E = E_1 \oplus E_2$, G decomposes as

$$G = \begin{pmatrix} G_1 & 0 \\ 0 & G_2 \end{pmatrix}, \quad (22)$$

where $G_1|_0 = \delta_0^n \otimes 1_{E_1}$, $G_2|_0 = 0$ and $\partial_0 G_2|_0 = \delta_0^n \otimes 1_{E_2}$.

We define the time dependent *Lagrange–Duhamel* vector field V_t on $\mathcal{E}_0^s(\mathbb{R})$ by: $\forall x \in M$,

$$V_t(\varphi)(x) := \int_{\mathbb{R}^n} d\vec{y} G(x^0 - t, \vec{x} - \vec{y}) N(\varphi, \partial\varphi)(t, \vec{y}) = \int_{y^0=t} d\vec{y} G_y(x) N(\varphi, \partial\varphi)(y), \quad (23)$$

where $G_y(x) := G(x - y)$. By Theorem 2.1 a map u is a solution of (1) iff

$$\frac{d(\Theta_t u)}{dt} + V_t(\Theta_t u) = 0, \quad (24)$$

The chronological exponential — The chronological exponential of $(V_t)_{t \in I}$ (if it exists) is the operator acting on \mathbb{F} defined by

$$\begin{aligned} T \exp \int_{t_1}^{t_2} d\tau V_\tau \cdot &:= \sum_{j=0}^{\infty} \int_{t_1 < \tau_1 < \dots < \tau_j < t_2} d\tau_1 \cdots d\tau_j (V_{\tau_j} \cdots V_{\tau_1} \cdot), \quad \text{if } t_2 > t_1 \\ \text{or} &:= \sum_{j=0}^{\infty} \int_{t_2 < \tau_j < \dots < \tau_1 < t_1} d\tau_1 \cdots d\tau_j (-1)^j (V_{\tau_j} \cdots V_{\tau_1} \cdot), \quad \text{if } t_2 < t_1, \end{aligned} \quad (25)$$

with the convention that the first term in the sum ($j = 0$) is the identity operator.

2 The Lagrange–Duhamel vector field formulation

The aim of this Section is to prove the following results.

Lemma 2.1 *Let J and I be two intervals of \mathbb{R} s.t. $J \subset I$. Assume that P_2 satisfies (9) and that either μ_0 in (9) is positive or I is bounded. Then the map*

$$\Theta : I \times \mathcal{F}^s(J) \longrightarrow \mathcal{E}_0^s(I)$$

defined by (13) exists and is continuous. Moreover there exists a constant $C_\Theta(I) > 0$ s.t.

$$\forall u \in \mathcal{F}^s(J), \forall t \in J, \quad \|\Theta_t u\|_{\mathcal{F}^s(I)} \leq C_\Theta(I) \|u\|_{\mathcal{F}^s(J)}.$$

Proposition 2.1 *Assume that P_2 satisfies (9) and that either μ_0 in (9) is positive or $I \subset \mathbb{R}$ is bounded. Then there exists some constant $Q_s > 0$ such that the following holds.*

Let $N : E_{(1)} \rightarrow E$ be a real analytic map of multiradius of convergence $\rho_{\otimes}(N) > 0$. Assume that: either

- (i) N_2 does not depend on ∂u and $s > n/2$; or*
- (ii) N_2 is affine in ∂u , i.e. (20) holds and $s > n/2 > s - r \geq 0$; or*
- (iii) $s > n/2 + 1$.*

Then $\exists \rho_{\otimes}(V)$ s.t. $Q_s \rho_{\otimes}(V) \geq \rho_{\otimes}(N)$ and $\forall (t, \varphi) \in I \times B_{\mathcal{E}_0^s(I)}(\rho_{\otimes}(V))$, the quantity

$$V(t, \varphi) = V_t(\varphi) := \int_{y^0=t} d\vec{y} G_y N(\varphi, \partial\varphi)(y)$$

is well-defined. Moreover the map $V : I \times B_{\mathcal{E}_0^s(\rho_{\otimes}(V))} \rightarrow \mathcal{E}_0^s(I)$ is continuous and $(V_t)_{t \in I}$ is a normal family of analytic maps of multiradius equal to $\rho_{\otimes}(V)$.

Theorem 2.1 *Assume the same hypotheses as in Proposition 2.1. Let $u \in \mathcal{F}^s(I)$ s.t. $\|u\|_{\mathcal{F}^s(I)} \leq \rho_{\otimes}(V)$. Then u belongs to $\mathcal{E}^s(I)$ (i.e. is a weak solution of $Lu + N(u, \partial u) = 0$) iff the map*

$$\begin{aligned} I &\longrightarrow \mathcal{E}_0^s(I) \\ t &\longmapsto \Theta_t u \end{aligned}$$

is \mathcal{C}^1 and satisfies (24), i.e. $\frac{d(\Theta_t u)}{dt} + V_t(\Theta_t u) = 0$.

Remark 2.1 *Analogues of Proposition 2.1 and Theorem 2.1 can be proved without difficulty in the case where $E = E_2$ (i.e. L is a fully second order operator), $1 = s \leq n/2$ and if $N = N_2$ is a polynomial of degree $\deg N \leq n/n - 2$. This is a consequence of the Sobolev embedding $H^1(\mathbb{R}^n) \hookrightarrow L^{2n/n-2}(\mathbb{R}^n)$, which allows to estimate the nonlinearity in $H^0(\mathbb{R}^n) = L^2(\mathbb{R}^n)$. The relevant cases are: $n = 2$ ($\deg N$ is arbitrary); $n = 3$ ($\deg N \leq 3$) and $n = 4$ ($\deg N \leq 2$). The proof is left to the Reader. The special case $n = 3$ and $N(u) = u^3$ will be treated in Section 3.*

2.1 Existence and continuity of Θ

For any $(\psi, \chi) \in \text{Cau}^s$ and $t \in \mathbb{R}$, we recall that $\Phi_t(\psi, \chi)$ is equal to the unique solution φ of $L\varphi = 0$ on \mathbb{R}^{n+1} s.t. $[\varphi]_t = (\psi, \chi)$. We also denote by $\Phi_t(\psi, \chi)$ the restriction of this map to any subset $I \times \mathbb{R}^n$. We set

$$\begin{aligned} \Phi : \mathbb{R} \times \text{Cau}^s &\longrightarrow \mathcal{E}_0^s(I) \\ (t, (\psi, \chi)) &\longmapsto \Phi_t(\psi, \chi). \end{aligned} \tag{26}$$

Proposition 2.2 *Assume that P_2 satisfies (9), then the linear map Φ defined by (26) is well-defined and continuous. Assume that either μ_0 in (9) is positive or $I \subset \mathbb{R}$ is bounded. Then there exists a constant $C_{\Phi}(I) > 0$ s.t.*

$$\|\Phi_t(\psi, \chi)\|_{\mathcal{F}^s(I)} \leq C_{\Phi}(I) \|(\psi, \chi)\|_{\text{Cau}^s}, \quad \forall (\psi, \chi) \in \text{Cau}^s. \tag{27}$$

Proof — Since Φ is linear in its second argument we can decompose the problem in two subcases and assume either $L = L_1 = \gamma^0 \partial_0 + P_1(\vec{\partial})$, or $L = L_2 = \partial_0^2 + P_2(\vec{\partial})$, separately.

Case $L = L_1$: we need to show that any solution φ to $\gamma^0 \partial_0 \varphi + P_1(\vec{\partial})\varphi = 0$, s.t. $\varphi|_t = \psi$ belongs to $\mathcal{F}^s(I)$ and depends continuously on (t, ψ) , where $\psi \in H^s(\mathbb{R}^n, E)$. Setting $\epsilon(\xi) := i(\gamma^0)^{-1} P_1(i\xi)$, the equation reads $\partial_0 \widehat{\varphi}(\tau, \xi) - i\epsilon(\xi) \widehat{\varphi}(\tau, \xi) = 0$, its solution is given by $\widehat{\varphi}(\tau, \xi) = e^{i\epsilon(\xi)(\tau-t)} \widehat{\psi}(\xi)$ and its time derivative by $\partial_0 \widehat{\varphi}(\tau, \xi) = i\epsilon(\xi) e^{i\epsilon(\xi)(\tau-t)} \widehat{\psi}(\xi)$. The result then follows by standard majorations and Lebesgue's dominated theorem.

Case $L = L_2$: We need to show that the solution φ of $(\partial_0)^2 \varphi + P_2(\vec{\partial})\varphi = 0$, s.t. $\varphi|_t = \psi$ and $\partial_0 \varphi|_t = \chi$ depends continuously on $(t, \psi, \chi) \in \mathbb{R} \times H^s(\mathbb{R}^n, E) \times H^{s-r}(\mathbb{R}^n, E)$. Assuming that $P_2(i\xi)$ is positive Hermitian and setting $\epsilon(\xi) := \sqrt{P_2(i\xi)}$, the equation reads $(\partial_0)^2 \widehat{\varphi}(\tau, \xi) + \epsilon(\xi)^2 \widehat{\varphi}(\tau, \xi) = 0$. Its solution is $\widehat{\varphi}(\tau, \xi) = \cos \epsilon(\xi)(\tau - t) \widehat{\psi}(\xi) + \epsilon(\xi)^{-1} \sin \epsilon(\xi)(\tau - t) \chi(\xi)$ and its time derivative is $\partial_0 \widehat{\varphi}(\tau, \xi) = -\epsilon(\xi) \sin \epsilon(\xi)(\tau - t) \widehat{\psi}(\xi) + \cos \epsilon(\xi)(\tau - t) \chi(\xi)$. The proof that $\varphi \in \mathcal{E}_0^s(I)$ and its continuous dependence on (t, ψ, χ) follows the same lines as for first order equations. However the factor $\epsilon(\xi)^{-1}$ in the expression of $\widehat{\varphi}$ may pose a slight difficulty in proving that φ is in $\mathcal{C}^0(I, H^s(\mathbb{R}^n, E))$ and that it depends continuously in t . In the ‘massive case’ (i.e. μ_0 in (9) is positive) this difficulty does not occur because of the inequality $|\epsilon(\xi)^{-1}| \leq (\alpha \mu_0)^{-1/2}$. In the ‘non massive’ case (i.e. μ_0 in (9) vanishes) we only have $|\epsilon(\xi)^{-1}| \leq \alpha^{-1/2} |\xi|^{-r}$. However by using the inequality:

$$\left| \frac{\sin \epsilon(\xi)t}{\epsilon(\xi)} \right| \leq \frac{\sqrt{1+t^2}}{\sqrt{1+\alpha|\xi|^{2r}}}, \quad (28)$$

we can prove the result by working with \mathcal{E}_0^s endowed with the norm

$$\|u\|_{\check{L}^\infty \text{Cau}^s} := \|u\|_{\check{L}^\infty(I, H^s)} + \|\partial_0 u\|_{L^\infty(I, H^{s-r})},$$

where $\|u\|_{\check{L}^\infty(I, H^s)} := \sup_{\tau \in I} \frac{\|u|_\tau\|_{H^s}}{\sqrt{1+\tau^2}}$. The conclusion follows if I is bounded, since then both norms $L^\infty \text{Cau}^s$ and $\check{L}^\infty \text{Cau}^s$ are equivalent. \square

Proof of Lemma 2.1 — Since the map Θ is obtained by composing Φ with the map $I \times \mathcal{F}^s(J) \longrightarrow I \times \text{Cau}^s$, $(t, u) \longmapsto (t, [u]_t)$, which is obviously continuous, Lemma 2.1 is a straightforward consequence of Proposition 2.2. \square

2.2 Estimate on the nonlinearity

The goal of this section is to collect results to prove Proposition 2.1. As a preliminary result we prove

Lemma 2.2 *Let \mathbb{X} and \mathbb{Y} be Banach spaces. Let $f = \sum_{p=0}^{\infty} f^{(p)}$ be a formal series from \mathbb{X} to \mathbb{Y} . Assume that its multiradius of convergence $\rho_\otimes(f) > 0$. Then the map f defined by $\forall \varphi \in \mathbb{X}$, $f(\varphi) = \sum_{p=0}^{\infty} f^{(p)}(\varphi)$ is \mathcal{C}^∞ on $B_\mathbb{X}(\rho_\otimes(f))$.*

In particular: $\forall r$ s.t. $0 < r < \rho_\otimes(f)$, $\forall \varphi, \psi \in \mathbb{X}$, $\forall h \in \mathbb{R}$ s.t. $\|\varphi\|_\mathbb{X}, \|\varphi + h\psi\|_\mathbb{X} \leq r$

$$\|f(\varphi + h\psi) - f(\varphi)\|_\mathbb{Y} \leq \frac{d[f]}{dz}(r) \|h\psi\|_\mathbb{X}. \quad (29)$$

and

$$\|f(\varphi + h\psi) - f(\varphi) - h\delta f_\varphi(\psi)\|_{\mathbb{Y}} \leq \frac{1}{2} \frac{d^2[f]}{dz^2}(r) h^2 \|\psi\|_{\mathbb{X}} \quad (30)$$

Proof — We prove only that f is \mathcal{C}^1 and hence (29) and (30) and leave the general case to the Reader. We first prove (29). Let $p \in \mathbb{N}$. From the identity $f^{(p)}(\varphi + h\psi) - f^{(p)}(\varphi) = \sum_{j=1}^p f_{\otimes}^{(p)}((\varphi + h\psi)^{\otimes j-1} \otimes h\psi \otimes \varphi^{\otimes p-j})$ we deduce

$$\|f^{(p)}(\varphi + h\psi) - f^{(p)}(\varphi)\|_{\mathbb{Y}} \leq \sum_{j=1}^p \|f_{\otimes}^{(p)}\|_{\otimes} \|\varphi + h\psi\|_{\mathbb{X}}^{j-1} \|h\psi\|_{\mathbb{X}} \|\varphi\|_{\mathbb{X}}^{p-j}.$$

Thus if $\|\varphi\|_{\mathbb{X}}, \|\varphi + h\psi\|_{\mathbb{X}} \leq r$,

$$\|f^{(p)}(\varphi + h\psi) - f^{(p)}(\varphi)\|_{\mathbb{Y}} \leq p \|f_{\otimes}^{(p)}\|_{\otimes} r^{p-1} \|h\psi\|_{\mathbb{X}}. \quad (31)$$

Hence by summing up on $p \in \mathbb{N}$ and using $\frac{d[f]}{dz} := \sum_{p=0}^{\infty} p \|f_{\otimes}^{(p)}\|_{\otimes} z^{p-1}$, we deduce that (29) holds if $r < \rho_{\otimes}(f)$.

The proof of (30) is similar. We start from the identity

$$f^{(p)}(\varphi + h\psi) - f^{(p)}(\varphi) - p f_{\otimes}^{(p)}(h\psi \otimes \varphi^{\otimes p-1}) = \sum_{j_1=1}^p \sum_{j_2=1}^{j_1-1} f_{\otimes}^{(p)}((\varphi + h\psi)^{\otimes j_2-1} \otimes (h\psi)^{\otimes 2} \otimes \varphi^{\otimes p-j_2-1}),$$

from which we deduce that, if $\|\varphi\|_{\mathbb{X}}, \|\varphi + h\psi\|_{\mathbb{X}} \leq r$,

$$\left\| f^{(p)}(\varphi + h\psi) - f^{(p)}(\varphi) - p f_{\otimes}^{(p)}(h\psi \otimes \varphi^{\otimes p-1}) \right\|_{\mathbb{Y}} \leq \frac{p(p-1)}{2} \|f_{\otimes}^{(p)}\|_{\otimes} r^{p-2} \|h\psi\|_{\mathbb{X}}^2.$$

Hence (30) follows by summing up on $p \in \mathbb{N}$. \square

Let V and W be two real vector spaces of (finite) dimension d_V and d_W respectively, let $k \in \mathbb{N}$ and $L \in \mathcal{Q}^k(V, W)$. Let $L_{\otimes} : V^{\otimes k} \rightarrow W$ the associated polarized linear map. Using bases on V and W , L_{\otimes} has the coordinates representation:

$$L_{\otimes}^i(z_1 \otimes \cdots \otimes z_k) := \sum_{j_1, \dots, j_k=1}^{d_V} L_{j_1, \dots, j_k}^i z_1^{j_1} \cdots z_k^{j_k} \quad \forall i = 1, \dots, d_W, \forall z_1, \dots, z_k \in V. \quad (32)$$

where, $\forall a$ s.t. $1 \leq a \leq k$, $(z_a^j)_{1 \leq j \leq d_V}$ are the coordinates of z_a and the coefficients L_{j_1, \dots, j_k}^i are symmetric in (j_1, \dots, j_k) . We set

$$|L| := \sum_{i=1}^{d_W} \sup_{1 \leq j_1, \dots, j_k \leq d_V} |L_{j_1, \dots, j_k}^i|. \quad (33)$$

One can easily check that (see (15))

$$\|L\|_{\otimes} \leq |L| \leq d_W \|L\|_{\otimes}. \quad (34)$$

The following result uses the fact that, if $s > n/2$, then $H^s(\mathbb{R}^n)$ is an algebra, i.e. the product of two functions $f, g \in H^s(\mathbb{R}^n)$ belongs to $H^s(\mathbb{R}^n)$ and there exists a constant Q_s s.t. $\|f\|_{H^s} \|g\|_{H^s} \leq Q_s \|fg\|_{H^s}$.

Lemma 2.3 *Let V and W be two real vector spaces of finite dimension, let $k \in \mathbb{N}$ and let $L \in \mathcal{Q}^k(V, W)$. Assume that $s > n/2$. Then one can define the homogeneous polynomial map $\mathcal{L} \in \mathcal{Q}^k((H^s(\mathbb{R}^n, V))^{\otimes k}, H^s(\mathbb{R}^n, W))$ by $\mathcal{L}_{\otimes}(\varphi_1 \otimes \cdots \otimes \varphi_k)(x) := L_{\otimes}(\varphi_1(x) \otimes \cdots \otimes \varphi_k(x))$ a.e., this map is linear continuous and satisfies*

$$\|\mathcal{L}\|^{\otimes} \leq Q_s^{k-1}|L|. \quad (35)$$

Proof — A straightforward recursion shows that $\|\varphi_1^{j_1} \cdots \varphi_k^{j_k}\|_{H^s} \leq Q_s^{k-1} \|\varphi_1^{j_1}\|_{H^s} \cdots \|\varphi_k^{j_k}\|_{H^s}$ and hence:

$$\begin{aligned} \|\mathcal{L}(\varphi_1 \otimes \cdots \otimes \varphi_k)\|_{H^s} &= \sum_{i=1}^{d_W} \|\mathcal{L}^i(\varphi_1 \otimes \cdots \otimes \varphi_k)\|_{H^s} \\ &\leq \sum_{i=1}^{d_W} \sum_{j_1, \dots, j_k=1}^{d_V} |L_{j_1, \dots, j_k}^i| \|\varphi_1^{j_1} \cdots \varphi_k^{j_k}\|_{H^s} \\ &\leq \sum_{i=1}^{d_W} \sup_{1 \leq j_1, \dots, j_k \leq d} |L_{j_1, \dots, j_k}^i| \sum_{j_1, \dots, j_k=1}^{d_V} Q_s^{k-1} \|\varphi_1^{j_1}\|_{H^s} \cdots \|\varphi_k^{j_k}\|_{H^s}. \end{aligned}$$

Hence the result follows by using (33). \square

In the following we use the notations:

$$\begin{aligned} \text{Cau}^s &\longrightarrow H^s(\mathbb{R}^n, E) \times H^{s-r}(\mathbb{R}^n, E_2) \times H^{s-1}(\mathbb{R}^n, \mathcal{L}(\mathbb{R}^n, E)) \\ \phi = (\varphi, \chi) &\longmapsto \phi^{(1)} = (\varphi, \chi, \partial_1 \varphi, \dots, \partial_n \varphi). \end{aligned} \quad (36)$$

As a first application of Lemma 2.3, given $N_1 = \sum_{k=0}^{\infty} N_1^{(k)} \in \mathbb{F}(E, E_1)$, we define for any $k \in \mathbb{N}$ the map $(\mathcal{N}_1^{(k)})_{\otimes} : (\text{Cau}^s)^{\otimes k} \longrightarrow H^s(\mathbb{R}^n, E_1)$ by $\forall (\phi_1, \dots, \phi_k) \in (\text{Cau}^s)^k$,

$$\mathcal{N}_1^{(k)}(\phi_1 \otimes \cdots \otimes \phi_k)(x) := N_1^{(k)}(\varphi_1(x) \otimes \cdots \otimes \varphi_k(x)), \quad \text{for a.e. } x \in \mathbb{R}^n. \quad (37)$$

We then deduce from (35) the estimate $\|(\mathcal{N}_1^{(k)})_{\otimes}\|^{\otimes} \leq Q_s^{k-1}|N_1^{(k)}|$. A similar estimate can be obtained for N_2 if this function does not depend on ∂u or if $s > n/2 + 1$.

However if N_2 is affine in ∂u , i.e. has the form (20) and if we suppose that $0 \leq s - r < n/2 < s$, then we use the fact that the product $(f, g) \longmapsto fg$ also maps continuously $H^s(\mathbb{R}^n) \times H^{s-r}(\mathbb{R}^n)$ to $H^{s-r}(\mathbb{R}^n)$ and that there exists a constant $q_{s,r}$ s.t.

$$\|fg\|_{H^{s-r}} \leq q_{s,r} Q_s \|f\|_{H^s} \|g\|_{H^{s-r}}. \quad (38)$$

This can be proved by splitting the product fg as the sum $T_f g + T_g f + R(f, g)$, where $(f, g) \longmapsto T_f g$ is the paraproduct and by estimating each term separately: $T_f g \in H^{s-r}(\mathbb{R}^n)$ because $f \in H^s(\mathbb{R}^n) \subset L^\infty(\mathbb{R}^n)$, $T_g f \in H^{s+(s-r)-n/2}(\mathbb{R}^n) \subset H^{s-r}(\mathbb{R}^n)$ because $s - r < n/2$ and $R(f, g) \in H^{s+(s-r)-n/2}(\mathbb{R}^n) \subset H^{s-r}(\mathbb{R}^n)$ because $s + (s - r) > n/2$ (see [3], Exercise A.5, page 109).

For the following remind the notation introduced in (36). We also use the notation $\delta_0 \phi := \chi$, $\delta_i \phi := \partial_i \varphi$, for $1 \leq i \leq n$, $\delta \phi := (\delta_\mu \phi)_{0 \leq \mu \leq n}$, $\forall \phi = (\varphi, \chi) \in \text{Cau}^s$.

Lemma 2.4 Let $N_2^{(k)} \in \mathcal{Q}^k(E_{(1)}, E_2)$ satisfying (20) with $J^{(k)} \in \mathcal{Q}^k(E, E_2)$ and $K_i^{(k)\mu} \in \mathcal{Q}^{k-1}(E_{(1)}, E_2)$. Assume that $0 \leq s-r < n/2 < s$. Then one can define the map $\mathcal{N}_2^{(k)}$ from $(\text{Cau}^s)^{\otimes k}$ to $H^{s-r}(\mathbb{R}^n, E_2)$ by $\mathcal{N}_2^{(k)}(\phi_1 \otimes \cdots \otimes \phi_k)(x) := N_2^{(k)}\left(\phi_1^{(1)}(x) \otimes \cdots \otimes \phi_k^{(1)}(x)\right)$, for a.e. $x \in \mathbb{R}^n$, this map is linear continuous and satisfies:

$$\|\mathcal{N}_2^{(k)}\|_{\otimes} \leq Q_s^{k-1} (|J^{(k)}| + q_{r,s}\sqrt{n}|K^{(k)}|) \quad \text{for} \quad |K^{(k)}| := \sup_{1 \leq i \leq \dim E} \sup_{0 \leq \mu \leq n} |K_i^{(k)\mu}|. \quad (39)$$

Proof — We start from the decomposition

$$\begin{aligned} \mathcal{N}_2^{(k)}(\phi_1 \otimes \cdots \otimes \phi_k) &= \mathcal{J}^{(k)}(\varphi_1 \otimes \cdots \otimes \varphi_k) \\ &\quad + \frac{1}{k} \sum_{a=1}^k \sum_{i=1}^{\dim E} \sum_{\mu=0}^n \mathcal{K}_i^{(k)\mu}(\varphi_1 \otimes \cdots \otimes \widehat{\varphi}_a \otimes \cdots \otimes \varphi_k) \delta_\mu \phi_a^i. \end{aligned} \quad (40)$$

Inequality (35) gives us $\|\mathcal{J}^{(k)}(\varphi_1 \otimes \cdots \otimes \varphi_k)\|_{H^s} \leq Q_s^{k-1}|J^{(k)}|\|\varphi_1\|_{H^s} \cdots \|\varphi_k\|_{H^s}$. This implies automatically a similar estimate in H^{s-r} . The H^{s-r} norm of the r.h.s. term in (40) is estimated by using (38):

$$\begin{aligned} &\leq \frac{q_{r,s}Q_s}{k} \sum_{a=1}^k \left(\sup_{1 \leq i \leq \dim E} \sup_{0 \leq \mu \leq n} \left\| \mathcal{K}_i^{(k)\mu}(\varphi_1 \otimes \cdots \otimes \widehat{\varphi}_a \otimes \cdots \otimes \varphi_k) \right\|_{H^s} \right) \left(\sum_{i=1}^{\dim E} \sum_{\mu=0}^n \|\delta_\mu \phi_a^i\|_{H^{s-r}} \right) \\ &\leq \frac{q_{r,s}Q_s}{k} \sum_{a=1}^k \left(|K^{(k)}| Q_s^{k-2} \|\varphi_1\|_{H^s} \cdots \|\widehat{\varphi}_a\|_{H^s} \cdots \|\varphi_k\|_{H^s} \right) \left(\sum_{\mu=0}^n \|\delta_\mu \phi_a\|_{H^{s-r}} \right), \end{aligned}$$

where we have used (35). However by using Cauchy–Schwarz inequality and the fact that $r \geq 1$ we have $\forall a = 1, \dots, k$,

$$\begin{aligned} \sum_{\mu=0}^n \|\delta_\mu \phi_a\|_{H^{s-r}} &\leq \|\chi_a\|_{H^{s-r}} + \sqrt{n} \left(\sum_{\mu=0}^n \|\partial_\mu \varphi_a\|_{H^{s-r}}^2 \right)^{1/2} \\ &\leq \|\chi_a\|_{H^{s-r}} + \sqrt{n} \|\varphi_a\|_{H^{s+1-r}} \leq \sqrt{n} \|\phi_a\|_{\text{Cau}^s}. \end{aligned}$$

Hence the H^{s-r} norm of the r.h.s. term in (40) is estimated by:

$$\begin{aligned} &\leq Q_s^{k-1} \frac{q_{r,s}\sqrt{n}}{k} |K^{(k)}| \sum_{a=1}^k \left(\|\varphi_1\|_{H^s} \cdots \|\widehat{\varphi}_a\|_{H^s} \cdots \|\varphi_k\|_{H^s} \right) \|\phi_a\|_{\text{Cau}^s} \\ &\leq Q_s^{k-1} q_{r,s}\sqrt{n} |K^{(k)}| \|\phi_1\|_{\text{Cau}^s} \cdots \|\phi_k\|_{\text{Cau}^s}. \end{aligned}$$

Hence (39) follows. \square

Let's summarize Lemmas 2.3 and 2.4. We can define for any $k \in \mathbb{N}$ the map $\mathcal{N}_{\otimes}^{(k)} : (\text{Cau}^s)^{\otimes k} \rightarrow \text{Cau}^s$ by: for a.e. $x \in \mathbb{R}^n$,

$$\mathcal{N}^{(k)}(\phi_1 \otimes \cdots \otimes \phi_k)(x) := \left(\iota \circ \mathcal{N}_1^{(k)}(\phi_1 \otimes \cdots \otimes \phi_k)(x), \mathcal{N}_2^{(k)}(\phi_1 \otimes \cdots \otimes \phi_k)(x) \right),$$

where $\iota : E_1 \rightarrow E$ is the natural inclusion. Remark that

$$\|\mathcal{N}^{(k)}(\phi_1 \otimes \cdots \otimes \phi_k)\|_{\text{Cau}^s} = \|\mathcal{N}_1^{(k)}(\varphi_1 \otimes \cdots \otimes \varphi_k)\|_{H^s} + \|\mathcal{N}_2^{(k)}(\phi_1 \otimes \cdots \otimes \phi_k)\|_{H^{s-r}}.$$

Proposition 2.3 *Assume that $N = \sum_{k=0}^{\infty} N^{(k)} \in \mathbb{F}(E_{(1)}, E)$ satisfies the same hypotheses as in Proposition 2.1. Then, for any $k \in \mathbb{N}$, the linear map $\mathcal{N}_{\otimes}^{(k)}$ from $(\text{Cau}^s)^{\otimes k}$ to Cau^s is continuous and satisfies :*

$$\|\mathcal{N}^{(k)}\|_{\otimes} \leq Q_s^k C(s, r, n) \|N^{(k)}\|_{\otimes}. \quad (41)$$

Hence if $\rho_{\otimes}(N) > 0$, then $\mathcal{N} = \sum_{k=0}^{\infty} \mathcal{N}^{(k)} \in \mathbb{F}_{Q_s^{-1}\rho_{\otimes}(N)}(\text{Cau}^s, \text{Cau}^s)$.

Proof — Case (i) where $s > n/2$ and $N_2^{(k)}$ does not depend on ∂u and Case (iii) where $s > n/2 + 1$ are similar and can be dealt by applying Lemma 2.3 for $L = N_1^{(k)}$ and $L = N_2^{(k)}$. We then obtain $\|\mathcal{N}^{(k)}\|_{\otimes} \leq Q_s^{k-1}(|N_1^{(k)}| + |N_2^{(k)}|)$. In Case (ii), we apply Lemma 2.3 for $L = N_1^{(k)}$ and Lemma 2.4 for $N_2^{(k)}$ to get $\|\mathcal{N}^{(k)}\|_{\otimes} \leq Q_s^{k-1}(|N_1^{(k)}| + |J^{(k)}| + q_{r,s}\sqrt{n}|K^{(k)}|)$. In any case (41) follows by applying (34) to $L = N_1^{(k)}, N_2^{(k)}, J^{(k)}, K^{(k-1)}$. As a consequence $\sum_{k=0}^{\infty} \|\mathcal{N}^{(k)}\|_{\otimes} R^k$ converges if $Q_s R < \rho_{\otimes}(N)$. \square

A first consequence of Proposition 2.3 is:

Proposition 2.4 *Let $s \in \mathbb{R}$ and $u \in \mathcal{F}^s(I)$. Assume that $Q_s \|u\|_{\mathcal{F}^s(I)} < \rho_{\otimes}(N)$. Assume that N satisfies the same hypotheses in Proposition 2.1. Then $\mathcal{N}(u, \partial u) \in \mathcal{C}^0(I, \text{Cau}^s)$, i.e. $\mathcal{N}_1(u) \in \mathcal{C}^0(I, H^s(\mathbb{R}^n, E))$ and $\mathcal{N}_2(u, \partial u) \in \mathcal{C}^0(I, H^{s-r}(\mathbb{R}^n, E_2))$.*

Proof — A straightforward consequence of Lemma 2.2, Proposition 2.3 and the continuity of $t \mapsto [u]_t \in \text{Cau}^s$. \square

2.3 The Lagrange–Duhamel vector field

We prove Proposition 2.1 — For any interval $I \subset \mathbb{R}$ and $k \in \mathbb{N}$, we define $V_{\otimes}^{(k)} : I \times (\mathcal{E}_0^s(I))^{\otimes k} \rightarrow \mathcal{E}_0^s(I)$ by

$$V^{(k)}(t, \varphi_1 \otimes \cdots \otimes \varphi_k) = V_t^{(k)}(\varphi_1 \otimes \cdots \otimes \varphi_k) := \Phi_t(\mathcal{N}^{(k)}([\varphi_1]_t \otimes \cdots \otimes [\varphi_k]_t))$$

$V_{\otimes}^{(k)}$ is continuous since it is the composition of the maps $I \times (\mathcal{E}_0^s(I))^{\otimes k} \ni (t, \varphi_1 \otimes \cdots \otimes \varphi_k) \mapsto (t, [\varphi_1]_t \otimes \cdots \otimes [\varphi_k]_t) \in I \times (\text{Cau}^s)^{\otimes k}$ (see (10)), $\mathcal{N}_{\otimes}^{(k)}$ (see Proposition 2.3) and Φ (see Proposition 2.2). By using (27) and (41) we deduce:

$$\|V^{(k)}\|_{\otimes} := \sup_{t \in I} \|V_t^{(k)}\|_{\otimes} \leq C_{\Phi}(I) \|\mathcal{N}^{(k)}\|_{\otimes} \leq C_{\Phi}(I) C(s, r, n) Q_s^k \|N^{(k)}\|_{\otimes}. \quad (42)$$

Setting $V_t := \sum_{k=0}^{\infty} V_t^{(k)}$, (42) implies that $(V_t)_{t \in I}$ is a normal family of analytic maps of multiradius of convergence $\rho_{\otimes}(V) \geq \rho_{\otimes}(N)/Q_s$.

To prove that $V := \sum_{k=0}^{\infty} V^{(k)}$ is continuous on $I \times B_{\mathcal{E}_0^s(I)}(0, \rho_{\otimes}(V))$, let $t, \tilde{t} \in I$, $\tilde{\varphi}, \varphi \in B_{\mathcal{E}_0^s(I)}(r)$, where $r < \rho_{\otimes}(V)$ and let us start from the inequality

$$\|V(\tilde{t}, \tilde{\varphi}) - V(t, \varphi)\|_{\mathcal{E}_0^s(I)} \leq \|\Phi_{\tilde{t}}(\mathcal{N}([\tilde{\varphi}]_{\tilde{t}}) - \mathcal{N}([\varphi]_t))\|_{\mathcal{E}_0^s(I)} + \|(\Phi_{\tilde{t}} - \Phi_t)(\mathcal{N}([\varphi]_t))\|_{\mathcal{E}_0^s(I)}. \quad (43)$$

Fix t, φ and $\varepsilon > 0$, then we deduce from Proposition 2.2 that, by choosing \tilde{t} sufficiently close to t , the last term in the r.h.s. of (43) is less than $\varepsilon/2$.

The first term in the r.h.s. of (43) can be estimated by using first (27) and second (29):

$$\begin{aligned} \|\Phi_{\tilde{t}}(\mathcal{N}([\tilde{\varphi}]_{\tilde{t}}) - \mathcal{N}([\varphi]_t))\|_{\mathcal{E}_0^s(I)} &\leq C_{\Phi}(I) \|\mathcal{N}([\tilde{\varphi}]_{\tilde{t}}) - \mathcal{N}([\varphi]_t)\|_{\text{Caus}} \\ &\leq C_{\Phi}(I) \frac{d[\mathcal{N}]}{dz}(r) \|[\tilde{\varphi}]_{\tilde{t}} - [\varphi]_t\|_{\text{Caus}}, \end{aligned}$$

and hence will also be smaller than $\varepsilon/2$ if we choose $|\tilde{t} - t|$ and $\|\tilde{\varphi} - \varphi\|_{\text{Caus}}$ sufficiently small.

To conclude observe that

$$V_t^{(k)}(\varphi_1 \otimes \cdots \otimes \varphi_k)(x) = \int_{y^0=t} d\vec{y} G(x-y) N^{(k)}((\varphi_1, \partial\varphi_1) \otimes \cdots \otimes (\varphi_k, \partial\varphi_k))(y).$$

This can be proven by using the properties of G (see (22)). Hence in particular

$$V_t^{(k)}(\varphi)(x) := \int_{\mathbb{R}^n} d\vec{y} G(x^0 - t, \vec{x} - \vec{y}) N^{(k)}(\varphi, \partial\varphi)(t, \vec{y}). \quad (44)$$

Formula (23) which is similar to (44) follows straightforwardly. \square

2.4 Derivability of Θ

We recall below Duhamel's formula (45). Recall that G is the distribution defined in (22). A generalization of Duhamel's formula for $L = L_2 = \square_g + m^2$ on a curved pseudo Riemannian manifold will given and proved in Section 3.

Proposition 2.5 *Let $f = (f_1, f_2)$ where $f_1 \in L_{loc}^1(\mathbb{R}, H^s(\mathbb{R}^n, E_1))$ and $f_2 \in L_{loc}^1(\mathbb{R}, H^{s-r}(\mathbb{R}^n, E_2))$. Assume that $u \in \mathcal{F}^s(I)$ is a solution of $Lu = f$. Then*

$$\forall x \in \mathbb{R}^{1+n}, \quad u(x) = \Theta_t u(x) + \int_t^{x^0} dy^0 \int_{\mathbb{R}^n} d\vec{y} G(x-y) f(y). \quad (45)$$

We are now in position to give the:

Proof of Theorem 2.1 — The key observation is that $[\Theta_t u]_t = [u]_t$ implies $N(\Theta_t u, \partial(\Theta_t u))|_t = N(u, \partial u)|_t$ and thus

$$V_t(\Theta_t u)(x) := \int_{\mathbb{R}^n} d\vec{y} G(x^0 - t, \vec{x} - \vec{y}) N(u, \partial u)(t, \vec{y}) \quad (46)$$

Now since u is a solution of $Lu + N(u, \partial u) = 0$, we deduce from Proposition 2.5 that $u(x) = \Theta_t u(x) - \int_t^{x^0} dy^0 \int_{\mathbb{R}^n} d\vec{y} G(x-y) N(u, \partial u)(y)$, which gives us thank to (46):

$$\forall x \in I \times \mathbb{R}^n, \quad u(x) = \Theta_t u(x) - \int_t^{x^0} dy^0 V_{y^0}(\Theta_{y^0} u)(x).$$

This implies the following identity $\forall t_1, t_2 \in I$:

$$\Theta_{t_2}u - \Theta_{t_1}u + \int_{t_1}^{t_2} dy^0 V_{y^0}(\Theta_{y^0}u) = 0 \quad \text{in } \mathcal{E}_0^s. \quad (47)$$

Lastly Lemma 2.1 and Proposition 2.1 imply that $I \ni t \mapsto V_t(\Theta_t u) \in \mathcal{E}_0^s(I)$ is continuous. Hence (47) implies that $I \ni t \mapsto \Theta_t u \in \mathcal{E}_0^s(I)$ is \mathcal{C}^1 and satisfies (24). \square

3 Curved space-times

We show here how Theorem 2.1 can be extended to field equations on a curved space-time. Let \mathcal{M} be smooth manifold equipped with a \mathcal{C}^∞ pseudo-Riemannian metric g of signature $(+, -, \dots, -)$. We denote by $\square_g = |g|^{-1} \partial_\mu (|g| g^{\mu\nu} \partial_\nu)$, where $|g| := \sqrt{|\det(g_{\mu\nu})|}$, the wave operator and set $L_g := \square_g + m^2$. A frame (e_0, \dots, e_n) is called *g-orthonormal* if $\langle e_\mu, e_\nu \rangle_g = \eta_{\mu\nu}$, where $\eta_{00} = 1$, $\eta_{ii} = -1$ if $i \neq 0$ and $\eta_{\mu\nu} = 0$ if $\mu \neq \nu$. We consider the non homogeneous scalar wave (or Klein–Gordon) equation on \mathcal{M} :

$$L_g u := |g|^{-1} \partial_\mu (|g| g^{\mu\nu} \partial_\nu u) + m^2 u = f. \quad (48)$$

Homogeneous (i.e. for $f = 0$) solutions u to (48) are the critical points of the action functional

$$\mathcal{A}(u) = \frac{1}{2} \int_{\mathcal{M}} [|\partial u|_g^2 + m^2 u^2] d\text{vol}_g,$$

where $d\text{vol}_g$ is the Riemannian volume element (in local coordinates x^μ , $d\text{vol}_g = |g| dx$) and $|\partial u|_g^2 := g^{\mu\nu} \partial_\mu u \partial_\nu u$. Similarly for any space-like hypersurface σ we let $d\mu_g$ denote the positive Riemannian measure on σ and N be the future oriented unit normal vector to σ . We then define

$$L^2(\sigma) := \{v : \sigma \rightarrow \mathbb{R} \text{ measurable s.t. } \|v\|_{L^2(\sigma)}^2 := \int_\sigma v^2 d\mu_g < +\infty\},$$

and, using a *g-orthonormal* frame (e_0, \dots, e_n) s.t. $e_0 = N$,

$$H_m^1(\sigma) := \{v : \sigma \rightarrow \mathbb{R} \text{ measurable s.t. } \int_\sigma (\sum_{i=0}^n \langle e_i, \nabla u \rangle_g^2 + m^2 u^2) d\mu_g < +\infty\}.$$

A hypersurface Σ is called *Cauchy* if any maximal smooth causal curve in \mathcal{M} intersects Σ at exactly one point (a smooth causal curve is a curve s.t. any vector which is tangent to it is time-like). If Σ_1 and Σ_2 are two space-like hypersurfaces, we write $\Sigma_1 \prec \Sigma_2$ if Σ_1 is in the past of Σ_2 and $\Sigma_1 \cap \Sigma_2 = \emptyset$. If u is a real valued map defined on a neighbourhood of Σ , we denote by $[u]_\Sigma = (u|_\Sigma, \langle N, \nabla u \rangle_g|_\Sigma)$ the Cauchy data of u along Σ . Our aim is to prove the existence and uniqueness of weak solutions to (48) with Cauchy conditions in $H_m^1(\Sigma) \times L^2(\Sigma)$ for some space-like Cauchy hypersurface Σ .

3.1 Existence of solutions to the linear problem

We first need generalizations of Proposition 2.2 to this context. Such results were proved by Y. Choquet-Bruhat, D. Chistodoulou and M. Francaviglia [13]. Here we present a more general version of their result in the case $s = r = 1$ by using the same techniques (see also the beautiful book by S. Alinhac [2]). We will make the following further hypotheses on (\mathcal{M}, g) : there exists a smooth ‘temporal function’ $\tau : \mathcal{M} \rightarrow \mathbb{R}$ and a smooth ‘radial function’ $\rho : \mathcal{M} \rightarrow [0, +\infty)$ and constants $A_1, A_2, A_3 > 0$ s.t.

$$|\nabla\tau|_g^2 > 0 \quad \text{everywhere;} \quad (49)$$

$$\forall t \in \mathbb{R}, \quad \Sigma_t := \tau^{-1}(t) \quad \text{is a space-like Cauchy hypersurface;} \quad (50)$$

$$\forall r > 0, \forall t_1, t_2 \in \mathbb{R} \text{ s.t. } t_1 < t_2, \quad \{x \in \mathcal{M}; \tau(x) \in [t_1, t_2], \rho(x) \leq r\} \text{ is compact.} \quad (51)$$

Moreover there exists some $R_0 > 0$, s.t., $\forall x \in \mathcal{M}$,

$$\rho(x) \geq R_0 \implies -A_3 \leq |\nabla\rho(x)|_g^2 < 0 \quad \text{and} \quad |\langle \nabla\rho, \nabla\tau(x) \rangle_g| \leq A_2/\rho(x); \quad (52)$$

$$\rho(x) \geq R_0 \implies A_1/\rho(x)^2 \leq |\nabla\tau(x)|_g^2. \quad (53)$$

Lastly define $\ell := |\nabla\tau|_g^{-1}$ (the *lapse function*) and $T := \ell\nabla\tau = \nabla\tau/|\nabla\tau|_g$. We assume that there exists a continuous function $B : \mathbb{R} \rightarrow [0, +\infty)$ s.t.

$$(n+1) (|\nabla T|_g^2 - 2|T^\mu\nabla_\mu T|_g^2)^{1/2} \leq (B \circ \tau)|\nabla\tau|_g \quad \text{on } \mathcal{M}. \quad (54)$$

Conditions (49) and (50) are equivalent to the assumption that (\mathcal{M}, g) is *globally hyperbolic* (see [6]). Conditions (53) (together with (52)) means that the lapse function grows at most linearly in ρ at spatial infinity. Condition (54) is an assumption on the curvature of the integral curves of the vector field T .

Given a smooth function $u : \mathcal{M} \rightarrow \mathbb{R}$ we define its *stress-energy tensor* $S(u)$ (associated with the action functional \mathcal{A} , see [23]), defined by

$$S_\nu^\mu(u) := g^{\mu\lambda}\partial_\lambda u \partial_\nu u - \frac{1}{2} (|\partial u|_g^2 - m^2 u^2) \delta_\nu^\mu.$$

We say that u has a *compact spatial support* if it vanishes on $\{x \in \mathcal{M}; \rho(x) \geq h(t)\}$ for some continuous function $h : \mathbb{R} \rightarrow [0, +\infty)$. If so and if σ is a space-like hypersurface (possibly with boundary), we define the energy

$$E_u(\sigma) := \int_\sigma \langle S(u)N, N \rangle_g d\mu_g.$$

Note that $E_u(\sigma)$ is always nonnegative. In particular if, on σ , we use a g -orthonormal frame (e_0, \dots, e_n) s.t. $e_0 = N$, then $\langle S(u)N, N \rangle_g = S_0^0(u) = \frac{1}{2} (\sum_{\mu=0}^n \langle e_\mu, \nabla u \rangle_g^2 + m^2 u^2)$.

For any interval $I \subset \mathbb{R}$, we define $\|u\|_{I,\tau} := \sup_{t \in I} E_u(\Sigma_t)^{1/2}$ and

$$\mathcal{F}_I^1(\Sigma_\tau) := \left\{ \begin{array}{l} \text{the closure of the set of smooth functions on } \tau^{-1}(I) \text{ with} \\ \text{compact spatial support in the topology induced by } \|\cdot\|_{I,\tau}. \end{array} \right.$$

We also define $\mathcal{F}_{loc}^1(\Sigma_\tau) := \{u : \mathcal{M} \rightarrow \mathbb{R}; \forall I \subset \mathbb{R} \text{ s.t. } I \text{ is bounded, } \|u\|_{I,\tau} < +\infty\}$. For any positive function $\beta : \mathbb{R} \rightarrow (0, +\infty)$ (actually we will use $\beta(t) = \exp\frac{1}{2} \int_0^t |B(s)| ds$), we set $\|u\|_{\beta,\tau} := \sup_{t \in \mathbb{R}} \beta(t)^{-1} E_u(\Sigma_t)^{1/2}$ and $\mathcal{F}_\beta^1(\Sigma_\tau) := \{u \in \mathcal{F}_{loc}^1(\Sigma_\tau); \|u\|_{\beta,\tau} < +\infty\}$. Lastly we set

$$L_{loc}^1(\mathbb{R}, L_\ell^2(\Sigma_\tau)) := \{f : \mathcal{M} \rightarrow \mathbb{R} \text{ measurable s.t. } [t \mapsto \|\ell f|_{\Sigma_t}\|_{L^2(\Sigma_t)}] \in L_{loc}^1(\mathbb{R})\}$$

and, for any interval $I \subset \mathbb{R}$, we note $\|f\|_{L^1(I, L_\ell^2(\Sigma_\tau))} := \int_I \|\ell f\|_{L^2(\Sigma_t)} dt$.

The existence result in [13] concerned weak solutions to (48) with a Cauchy data on a hypersurface Σ_t . The following result extends this with the notable difference that we allow more general Cauchy hypersurfaces. Fixing τ (and hence the foliation $(\Sigma_t)_t$) we say that a space-like hypersurface $\widehat{\Sigma}$ is admissible if it is a Cauchy hypersurface and if: (i) $\exists t_1, t_2 \in \mathbb{R}$ s.t. $t_1 < t_2$ and $\Sigma_{t_1} \prec \widehat{\Sigma} \prec \Sigma_{t_2}$; (ii) if \widehat{N} denotes the future oriented normal to $\widehat{\Sigma}$, $C(\widehat{\Sigma}) := \sup_{\widehat{\Sigma}} \langle \widehat{N}, T \rangle_g < +\infty$.

Theorem 3.1 *Assume that (\mathcal{M}, g) satisfies Hypotheses (49)–(54). Let $\widehat{\Sigma} \subset \mathcal{M}$ be a space-like admissible hypersurface and let $t_1, t_2 \in \mathbb{R}$ s.t. $\Sigma_{t_1} \prec \widehat{\Sigma} \prec \Sigma_{t_2}$. Then for any $(u_0, u_1) \in H_m^1(\widehat{\Sigma}) \times L^2(\widehat{\Sigma})$ and $f \in L_{loc}^1(\mathbb{R}, L_\ell^2(\Sigma_\tau))$, there exists a unique weak solution $u \in \mathcal{F}_{loc}^1(\Sigma_\tau)$ to (48) s.t. $[u]_{\widehat{\Sigma}} = (u_0, u_1)$. Moreover for $t = t_1$ or t_2 ,*

$$E_u(\Sigma_t)^{1/2} \leq \sqrt{2}\beta(t_1, t_2) \left(C(\widehat{\Sigma}) E_u(\widehat{\Sigma}) \right)^{1/2} + \sqrt{2}\beta(t_1, t_2)^2 \|f\|_{L^1([t_1, t_2], L_\ell^2(\Sigma_\tau))}, \quad (55)$$

where $\beta(t_1, t_2) := e^{\frac{1}{2} \int_{t_1}^{t_2} |B(s)| ds}$ and $C(\widehat{\Sigma}) := \sup_{\widehat{\Sigma}} \langle \widehat{N}, T \rangle_g$.

Proof — The main point is to obtain the a priori estimate (55) for any solution u to (48). Without loss of generality we will content ourself to prove that

$$E_u(\Sigma_{t_2})^{1/2} \leq \sqrt{2} e^{\frac{1}{2} \int_{t_1}^{t_2} |B(s)| ds} \left(C(\widehat{\Sigma}) E_u(\widehat{\Sigma}) \right)^{1/2} + \sqrt{2} e^{\int_{t_1}^{t_2} |B(s)| ds} \int_{t_1}^{t_2} \|\ell f|_{\Sigma_s}\|_{L^2} ds. \quad (56)$$

Step 1: Use of conservation law — Consider a compact domain $D \subset \tau^{-1}([t_1, t_2])$, the boundary ∂D of which is composed of three smooth components


$$\partial D = (\Sigma_{t_2} \cap \overline{D}) + \Lambda - (\Sigma_{t_1} \cap \overline{D}),$$

where the signs give the orientation. We assume that Λ is space-like and that the normal vector N to it is future-pointing, hence $(\Sigma_{t_2} \cap \overline{D}) \cup \Lambda$ forms the top of D , whereas $\Sigma_{t_1} \cap \overline{D}$ is the bottom (see the end of the proof for the construction of D).

Let $\theta : \mathcal{M} \rightarrow \mathbb{R}$ be the function which coincides with τ on $\widehat{\Sigma}$ and which is invariant by the flow of T and, for $t \in [t_1, t_2]$, consider the domain

$$D_t := \{x \in D; \theta(x) < \tau(x) < t\}$$

(points in D_t are points of D which are in the future of $\widehat{\Sigma}$ and in the past of Σ_t , see the figure). Note that $\partial D_t = (\Sigma_t \cap \overline{D_t}) + (\Lambda \cap \overline{D_t}) - \widehat{\Sigma}_{<t}$, where $\Sigma_t \cap \overline{D_t} = \{x \in \overline{D}; \theta(x) <$



$\tau(x) = t$ and $\widehat{\Sigma}_{<t} = \{x \in \overline{D}; \theta(x) = \tau(x) < t\}$.

Let us apply Stokes theorem to $S(u)T$ on D_t . We get (writing $S = S(u)$ for shortness):

$$\int_{D_t} \nabla_\mu (S_\nu^\mu T^\nu) d\text{vol}_g = \int_{\partial D_t} \langle N, ST \rangle_g d\mu_g. \quad (57)$$

Since u is a solution of (48), the stress-energy tensor satisfies the relation $\nabla_\mu S_\nu^\mu = f \partial_\nu u$, see e.g. [23]. Hence the l.h.s. of (57) reads

$$\int_{D_t} \nabla_\mu (S_\nu^\mu T^\nu) d\text{vol}_g = \int_{D_t} f(\partial_\nu u) T^\nu d\text{vol}_g + \int_{D_t} S_\nu^\mu \nabla_\mu T^\nu d\text{vol}_g. \quad (58)$$

(i) *Estimation of the first term in the r.h.s of (58)* — Using the coarea formula, we get

$$\left| \int_{D_t} f T^\nu \partial_\nu u d\text{vol}_g \right| \leq \int_{D_t} |f| |T^\nu \partial_\nu u| d\text{vol}_g = \int_{t_1}^t ds \int_{\Sigma_s \cap \overline{D_t}} |T^\nu \partial_\nu u| |f| \frac{d\mu_g}{|\nabla \tau|_g}.$$

Since T coincides with the normal vector N to Σ_s , we have: $|T^\nu \partial_\nu u| = |\langle N, \nabla u \rangle_g| \leq \sqrt{2\langle SN, N \rangle_g}$. Hence by Cauchy–Schwarz

$$\begin{aligned} \left| \int_{D_t} f T^\nu \partial_\nu u d\text{vol}_g \right| &\leq \int_{t_1}^t ds \left(\int_{\Sigma_s \cap \overline{D_t}} 2\langle SN, N \rangle_g d\mu_g \right)^{1/2} \left(\int_{\Sigma_s \cap \overline{D_t}} (\ell f)^2 d\mu_g \right)^{1/2} \\ &= \int_{t_1}^t ds \sqrt{2E_u(\Sigma_s \cap \overline{D_t})} \|\ell f|_{\Sigma_s \cap \overline{D_t}}\|_{L^2}. \end{aligned}$$

(ii) *Estimation of the second term in the r.h.s of (58)* — The Cauchy–Schwarz inequality gives us:

$$|S_\nu^\mu \nabla_\mu T^\nu|^2 \leq \left(\sum_{\mu, \nu=0}^n (S_\nu^\mu)^2 \right) \left(\sum_{\mu, \nu=0}^n (\nabla_\mu T^\nu)^2 \right),$$

however a difficulty is that the r.h.s. of this inequality depends on the choice of the frame (e_0, \dots, e_n) used in the decomposition of the tensors S and ∇T . We choose a g -orthonormal frame (e_0, \dots, e_n) s.t. $e_0 = T$. Observe then that $|S_\nu^\mu| \leq S_0^0, \forall \mu, \nu$ and

thus⁴

$$\sum_{\mu,\nu=0}^n (S_\nu^\mu)^2 \leq (n+1)^2 (S_0^0)^2 = (n+1)^2 \langle T, ST \rangle_g^2.$$

Next let us introduce the tensor $h_{\mu\nu} := 2T_\mu T_\nu - g_{\mu\nu}$. We note that in the previously chosen g -orthonormal frame we have $h_{\mu\nu} = \delta_{\mu\nu}$ and hence that

$$\sum_{\mu,\nu=0}^n (\nabla_\mu T^\nu)^2 = \nabla_\mu T^\nu \nabla_\lambda T^\sigma h_{\nu\sigma} h^{\mu\lambda} =: |\nabla T|_h^2.$$

We thus deduce that

$$|S_\nu^\mu \nabla_\mu T^\nu| \leq (n+1) \langle T, ST \rangle_g |\nabla T|_h,$$

where the r.h.s. is now frame independent. Lastly a computation (using $|T|_g^2 = 1$, which implies $g_{\lambda\nu} T^\nu \nabla_\mu T^\lambda = 0$) shows that $|\nabla T|_h^2 = |\nabla T|_g^2 - 2|T^\mu \nabla_\mu T|_g^2$. Hence $|S_\nu^\mu \nabla_\mu T^\nu| \leq (n+1)(|\nabla T|_g^2 - 2|T^\mu \nabla_\mu T|_g^2)^{1/2} S_0^0$ and using the fact that T coincides with the future pointing normal vector to Σ_s , (54) and the coarea formula,

$$\begin{aligned} \left| \int_{D_t} S_\nu^\mu \nabla_\mu T^\nu d\text{vol}_g \right| &\leq \int_{D_t} B(\tau) |\nabla \tau|_g S_0^0 d\text{vol}_g \\ &= \int_{t_1}^t B(s) ds \int_{\Sigma_s \cap \overline{D_t}} \langle ST, T \rangle_g d\mu_g = \int_{t_1}^t B(s) E_u(\Sigma_s \cap \overline{D_t}) ds. \end{aligned}$$

Summarizing with the previous step we deduce from (58)

$$\left| \int_{D_t} \nabla_\mu (S_\nu^\mu T^\nu) d\text{vol}_g \right| \leq \int_{t_1}^t \|\ell f|_{\Sigma_s}\|_{L^2} \sqrt{2E_u(\Sigma_s \cap \overline{D_t})} ds + \int_{t_1}^t B(s) E_u(\Sigma_s \cap \overline{D_t}) ds,$$

which, in view of (57) gives:

$$\int_{\partial D_t} \langle N, ST \rangle_g d\mu_g \leq \int_{t_1}^t \|\ell f|_{\Sigma_s}\|_{L^2} \sqrt{2E_u(\Sigma_s \cap \overline{D_t})} ds + \int_{t_1}^t B(s) E_u(\Sigma_s \cap \overline{D_t}) ds. \quad (59)$$

(iii) *Lower estimation of the l.h.s. of (59)* — Using the fact that $T = N$ on Σ_t and denoting by \widehat{N} the future pointing normal to $\widehat{\Sigma}$, we decompose

$$\int_{\partial D_t} \langle N, ST \rangle_g d\mu_g = E_u(\Sigma_t \cap \overline{D_t}) + \int_{D_t \cap \Lambda} \langle ST, N \rangle_g d\mu_g - \int_{\widehat{\Sigma}_{<t}} \langle ST, \widehat{N} \rangle_g d\mu_g.$$

However $\langle ST, N \rangle_g \geq 0$ on $D_t \cap \Lambda$. This follows from $|N|_g^2 = 1$, $N^0 > 0$ and from the following identity, valid in a g -orthonormal frame (e_0, \dots, e_n) s.t. $e_0 = T$:

$$2N^0 \langle ST, N \rangle_g = |N|_g^2 (u^0)^2 + \sum_{i=1}^n (N^0 u^i - N^i u^0)^2 + (N^0)^2 m^2 u^2, \quad (60)$$

⁴This inequality is true for any vector valued field u . Actually using the fact that u is a scalar field one can get the improved inequality $\sum_{\mu,\nu=0}^n (S_\nu^\mu)^2 \leq (n+3)(S_0^0)^2$.

where $N^\mu = \langle e_\mu, N \rangle_g$ and $u^\mu := \langle e_\mu, \nabla u \rangle_g$. Hence

$$E_u(\Sigma_t \cap \overline{D_t}) - \int_{\widehat{\Sigma}_{<t}} \langle ST, \widehat{N} \rangle_g d\mu_g \leq \int_{\partial D_t} \langle N, ST \rangle_g d\mu_g. \quad (61)$$

(iv) *Conclusion* — For any $s \in [t_1, t]$ let $\widehat{\Sigma}_{\geq s} := \{x \in \widehat{\Sigma}; \tau(x) \geq s\}$ and set

$$e(s) := E_u(\Sigma_s \cap \overline{D_t}) + \int_{\widehat{\Sigma}_{\geq s}} \langle ST, \widehat{N} \rangle_g d\mu_g.$$

We will prove that the l.h.s. of (61) is equal to $e(t) - e(t_1)$. Observe that, because of $\Sigma_{t_1} \cap \overline{D_t} = \emptyset$ and $\widehat{\Sigma}_{\geq t_1} = \widehat{\Sigma} \cap D$, $e(t_1) = \int_{\widehat{\Sigma} \cap D} \langle ST, \widehat{N} \rangle_g d\mu_g$. But, since $\widehat{\Sigma}_{\geq t} \cap \widehat{\Sigma}_{<t} = \emptyset$ and $\widehat{\Sigma}_{\geq t} \cup \widehat{\Sigma}_{<t} = \widehat{\Sigma} \cap D$, the latter decomposes as:

$$e(t_1) = \int_{\widehat{\Sigma} \cap D} \langle ST, \widehat{N} \rangle_g d\mu_g = \int_{\widehat{\Sigma}_{<t}} \langle ST, \widehat{N} \rangle_g d\mu_g + \int_{\widehat{\Sigma}_{\geq t}} \langle ST, \widehat{N} \rangle_g d\mu_g.$$

Hence $e(t) - e(t_1) = E_u(\Sigma_t \cap \overline{D_t}) - \int_{\widehat{\Sigma}_{<t}} \langle ST, \widehat{N} \rangle_g d\mu_g$, so that (61) reads $e(t) - e(t_1) \leq \int_{\partial D_t} \langle N, ST \rangle_g d\mu_g$. By using (59) and the fact that $E_u(\Sigma_s \cap \overline{D_t}) \leq e(s)$ we deduce (setting $F(s) := \sqrt{2} \|\ell f|_{\Sigma_s}\|_{L^2}$) that:

$$e(t) - e(t_1) \leq \int_{t_1}^t ds F(s) \sqrt{e(s)} + \int_{t_1}^t B(s) e(s) ds. \quad (62)$$

Step 3: Using Gronwall lemma — Set $K := e(t_1) + \int_{t_1}^t ds F(s) \sqrt{e(s)}$. Then (62) (by replacing t by t') implies easily $e(t') \leq K + \int_{t_1}^{t'} B(s) e(s) ds$, $\forall t' \in [t_1, t]$. Using Gronwall Lemma we deduce that

$$e(t') \leq K e^{\int_{t_1}^{t'} B(s) ds}, \quad \forall t' \in [t_1, t].$$

Replacing K by its value, setting $\psi(t) := \sup_{t_1 \leq s \leq t} \sqrt{e(s)}$ and taking the supremum over $t' \in [t_1, t]$, we obtain

$$\psi(t)^2 \leq \left(\psi(t_1)^2 + \int_{t_1}^t ds F(s) \psi(s) \right) e^{\int_{t_1}^t B(s) ds} \leq \left(\psi(t_1)^2 + \psi(t) \int_{t_1}^t F(s) ds \right) e^{\int_{t_1}^t B(s) ds},$$

which implies $\psi(t) \leq \psi(t_1) e^{\frac{1}{2} \int_0^t B(s) ds} + \left(\int_{t_1}^t F(s) ds \right) e^{\int_{t_1}^t B(s) ds}$. Applying this for $t = t_2$ and using $e(t_2) = E_u(\Sigma_{t_2} \cap \overline{D})$, we get

$$E_u(\Sigma_{t_2} \cap \overline{D})^{1/2} \leq e^{\frac{1}{2} \int_{t_1}^{t_2} B(s) ds} e(t_1)^{1/2} + e^{\int_{t_1}^{t_2} B(s) ds} \int_{t_1}^{t_2} F(s) ds. \quad (63)$$

Step 4: Controlling $e(t_1)$ by $E_u(\widehat{\Sigma})$ — Using an identity similar to (60) (where T is replaced by \widehat{N} , N is replaced by T and we use a g -orthonormal frame $(\widehat{e}_0, \dots, \widehat{e}_n)$ s.t. $\widehat{e}_0 = \widehat{N}$) we prove that

$$\langle ST, \widehat{N} \rangle_g \leq 2 \langle T, \widehat{N} \rangle_g \left(\frac{1}{2} \sum_{\mu=0}^n \langle \widehat{e}_\mu, \nabla u \rangle_g^2 + \frac{1}{2} m^2 u^2 \right) = 2 \langle T, \widehat{N} \rangle_g \langle S \widehat{N}, \widehat{N} \rangle_g.$$

This hence implies that

$$e(t_1) = \int_{\widehat{\Sigma} \cap D} \langle ST, \widehat{N} \rangle_g d\mu_g \leq 2 \sup_{\widehat{\Sigma}} \langle T, \widehat{N} \rangle_g E_u(\widehat{\Sigma} \cap \overline{D}) \leq 2C(\widehat{\Sigma}) E_u(\widehat{\Sigma}).$$

Thus we deduce from (63)

$$E_u(\Sigma_{t_2} \cap \overline{D})^{1/2} \leq \sqrt{2} e^{\frac{1}{2} \int_{t_1}^{t_2} B(s) ds} \left(C(\widehat{\Sigma}) E_u(\widehat{\Sigma}) \right)^{1/2} + e^{\int_{t_1}^{t_2} B(s) ds} \int_{t_1}^{t_2} F(s) ds. \quad (64)$$

Step 5: Global estimate — Now, for any $R > 0$, set $K_R := \{x \in \mathcal{M}; \rho(x) \leq R\}$. In order to obtain (56) it suffices to prove that there exists some $R_0 > 0$ s.t., for any $R > R_0$, there exists a domain D satisfying the previous properties and s.t. $\Sigma_{t_2} \cap K_R \subset \Sigma_{t_2} \cap \overline{D}$. Indeed if so we deduce from (64)

$$E_u(\Sigma_{t_2} \cap K_R)^{1/2} \leq E_u(\Sigma_{t_2} \cap \overline{D})^{1/2} \leq \sqrt{2} e^{\frac{1}{2} \int_{t_1}^{t_2} B(s) ds} \left(C(\widehat{\Sigma}) E_u(\widehat{\Sigma}) \right)^{1/2} + e^{\int_{t_1}^{t_2} B(s) ds} \int_{t_1}^{t_2} F(s) ds$$

Since this inequality holds for any $R > 0$, it thus implies (56).

Step 6: Construction of D — Here we need Hypotheses (51) to (53). Set $t := t_2 - t_1$ and assume that $t > 0$. For any $R > R_0$ we will construct a smooth function $\tilde{\tau} : \mathcal{M} \rightarrow \mathbb{R}$ and find some $\overline{R} > R$ s.t.

- (i) $\forall x \in K_R, \tilde{\tau}(x) = \tau(x) - t = \tau(x) - t_2 + t_1$;
- (ii) $\forall x \notin K_{\overline{R}}, \tilde{\tau}(x) = \tau(x)$;
- (iii) $|\nabla \tilde{\tau}|_g^2 > 0$ everywhere, in particular the level sets of $\tilde{\tau}$ are space-like hypersurfaces.

If so $D := \{x \in \mathcal{M}; \tau(x) > t_1, \tilde{\tau}(x) < t_1\}$ satisfies all the previously required properties. To construct $\tilde{\tau}$, we set $\tilde{\tau} = \tau - t\chi \circ \rho$, where $\chi \in \mathcal{C}^0([0, +\infty), [0, 1])$ is piecewise \mathcal{C}^∞ and has to be suitably chosen. Conditions (i) and (ii) translate respectively as: (i)' $\forall r \leq R, \chi(r) = 1$; (ii)' $\forall r \geq \overline{R}, \chi(r) = 0$. A simple computation using (52) and (53) shows that Condition (iii) is satisfied if

$$A_3 t^2 r^2 (\chi'(r))^2 + 2A_2 t r |\chi'(r)| < A_1.$$

This condition is fulfilled if we choose $\alpha > 0$ s.t. $A_3 \alpha^2 + 2A_2 \alpha < A_1$, $\overline{R} = R e^{t/\alpha}$ and set $\chi(r) = 1 - \frac{\alpha}{t} \log \frac{r}{\overline{R}}, \forall r \in [R, \overline{R}]$ (all that works because $\int_R^\infty \frac{dr}{r} = +\infty$).

Step 7: Conclusion — Thanks to the works of J. Hadamard, M. Riesz and the results by J. Leray [28], one can construct fundamental solutions for the operator L and solve the Cauchy problem for smooth Cauchy data (see [17, 6]). By using the density of smooth compactly supported functions in $L^2(\Sigma_{t_1})$ and $H_m^1(\Sigma_{t_1})$ and (55), we deduce the existence. The uniqueness is a straightforward consequence of (55). \square

Note that similar results exist for higher (integer) order Sobolev spaces and for Cauchy data on a hypersurface which belongs to the family $(\Sigma_t)_t$, see [13] and also [2]. Theorem 3.1 has the following consequence which is a substitute for Proposition 2.2. Set $\mathcal{E}_{0,\beta}^1(\Sigma_\tau) := \{\varphi \in \mathcal{F}_\beta^1(\Sigma_\tau); \square_g \varphi + m^2 \varphi = 0\}$.

Corollary 3.1 *Let (\mathcal{M}, g) be a n -dimensional Lorentzian manifold. Assume that there exist functions $\tau, \rho \in \mathcal{C}^\infty(\mathcal{M}, \mathbb{R})$ which satisfy (49)–(54). Set $\beta(t) = e^{\frac{1}{2}| \int_0^t B(s) ds |}$. Then for any admissible hypersurface σ , there exists a continuous linear map*

$$\begin{aligned} \Phi_\sigma : H_m^1(\sigma) \times L^2(\sigma) &\longrightarrow \mathcal{E}_{0,\beta}^1(\Sigma_\tau) \\ (\psi, \chi) &\longmapsto \Phi_\sigma(\psi, \chi), \end{aligned}$$

where $\Phi_\sigma(\psi, \chi)$ is equal to the unique solution φ to $L_g \varphi = \square_g \varphi + m^2 \varphi = 0$ with the Cauchy data $[\varphi]_\sigma = (\psi, \chi)$.

Thanks to this result we can define for any admissible hypersurface σ the continuous map

$$\Theta_\sigma : \mathcal{F}_\beta^1(\Sigma_\tau) \longrightarrow \mathcal{E}_{0,\beta}^1(\Sigma_\tau)$$

defined by $\Theta_\sigma(u) := \Phi_\sigma([u]_\sigma)$. The following result will also be useful.

Lemma 3.1 *Let $f \in L_{loc}^1(\mathbb{R}, L_\ell^2(\Sigma_\tau))$ and $u \in \mathcal{F}_{loc}^1(\Sigma_\tau)$ be a solution of $\square_g u + m^2 u = f$. Let $\widehat{\Sigma}$ be an admissible hypersurface s.t. $\Sigma_{t_1} \prec \widehat{\Sigma} \prec \Sigma_{t_2}$. Then*

$$E_u(\widehat{\Sigma}) \leq 2C(\widehat{\Sigma}) \left[(1 + \|B\|_{L^1([t_1, t_2])}) \|u\|_{[t_1, t_2], \tau} + \sqrt{2} \|f\|_{L^1([t_1, t_2], L_\ell^2)} \|u\|_{[t_1, t_2], \tau}^{1/2} \right]. \quad (65)$$

Sketch of the proof — The proof is based on the same techniques as in the proof of Theorem 3.1: one starts from the identity $\int_\Delta \nabla_\mu (S_\nu^\mu T^\nu) d\text{vol}_g = \int_{\partial\Delta} \langle ST, N \rangle_g d\mu_g$, with the same vector field T . The difference is the domain of integration which is now $\Delta := \{x \in D; t_1 < \tau(x) < \theta(x)\}$. Also the reasoning is simpler, for we already know that $\|u\|_{[t_1, t_2], \tau}$ is bounded and hence we do not need to use Gronwall lemma. This leads to

$$\int_{\widehat{\Sigma}} \langle ST, \widehat{N} \rangle_g d\mu_g \leq (1 + \|B\|_{L^1([t_1, t_2])}) \|u\|_{[t_1, t_2], \tau} + \sqrt{2} \|f\|_{L^1([t_1, t_2], L_\ell^2)} \|u\|_{[t_1, t_2], \tau}^{1/2}.$$

Estimate (65) follows then from the inequality $\langle S\widehat{N}, \widehat{N} \rangle_g \leq 2\langle T, \widehat{N} \rangle_g \langle ST, \widehat{N} \rangle_g$, which implies $E_u(\widehat{\Sigma}) \leq 2 \sup_{\widehat{\Sigma}} \langle T, \widehat{N} \rangle_g \int_{\widehat{\Sigma}} \langle ST, \widehat{N} \rangle_g d\mu_g$. \square

3.2 A generalization of Duhamel's formula

Our aim is here to prove a ‘curved’ version of Duhamel’s formula. Beside the foliation of \mathcal{M} by the level sets $\Sigma_t := \tau^{-1}(t)$, we also consider a family $(\sigma_s)_{s \in \mathbb{R}}$ of admissible Cauchy space-like hypersurfaces, *which may not form a foliation of \mathcal{M} in general*. We assume that there exists an n -dimensional manifold $\underline{\sigma}$ (the model for each σ_s) and a map $F \in \mathcal{C}^\infty(\mathbb{R} \times \underline{\sigma}, \mathcal{M})$ s.t. for any $s \in \mathbb{R}$, $F_s := F(s, \cdot)$ is an embedding of $\underline{\sigma}$, the image of which is σ_s . On each σ_s we define the function $\lambda_s \in \mathcal{C}^\infty(\sigma_s, \mathbb{R})$ by $\lambda_s \circ F_s := \langle \frac{\partial F}{\partial s}(s, \cdot), N_s \circ F_s \rangle_g$, where N_s is

the future pointing normal vector to σ_s . We call $(\sigma_s)_{s \in \mathbb{R}}$ a *smooth family of admissible Cauchy hypersurfaces*.

For any $s \in \mathbb{R}$, we denote by $\{x \succ \sigma_s\}$ (resp. $\{x \prec \sigma_s\}$) the subset of $\mathcal{M} \setminus \sigma_s$ which are in the future (resp. the past) of σ_s , similarly $\{x \succcurlyeq \sigma_s\} := \sigma_s \cup \{x \succ \sigma_s\}$ ($\{x \preccurlyeq \sigma_s\} := \sigma_s \cup \{x \prec \sigma_s\}$). We let $Y_{\sigma_s} \in L^\infty(\mathcal{M})$ be s.t. $Y_{\sigma_s} = 1$ on $\{x \succcurlyeq \sigma_s\}$ and $Y_{\sigma_s} = 0$ on $\{x \prec \sigma_s\}$.

We let $f \in L^1_{loc}(\mathbb{R}, L^2_\ell(\Sigma_\tau))$ and we assume that, for a.e. $s \in \mathbb{R}$, $\lambda_s f|_{\sigma_s} \in L^2(\sigma_s)$ and $[s \mapsto \|\lambda_s f|_{\sigma_s}\|_{L^2}]$ belongs to $L^1_{loc}(\mathbb{R})$. We then define:

$$\gamma_s f : \quad \text{the unique solution of} \quad \begin{cases} \gamma_s f = 0 & \text{on } \{x \prec \sigma_s\} \\ \gamma_s f = \Phi_{\sigma_s}(0, \lambda_s f|_{\sigma_s}) & \text{on } \{x \succcurlyeq \sigma_s\}, \end{cases}$$

$$\Gamma_s f : \quad \text{the unique solution of} \quad \begin{cases} \Gamma_s f = 0 & \text{on } \{x \prec \sigma_s\} \\ L_g(\Gamma_s f) = f Y_{\sigma_s} & \text{on } \mathcal{M}. \end{cases}$$

For any $y \in \mathcal{M}$ we let G_y be the solution of $L_g G_y = 0$ with the Cauchy data $G_y|_\sigma = 0$ and $\langle N, \nabla G_y \rangle_g|_\sigma = \delta_y$, where σ is a Cauchy hypersurface which contains y . Then, still if $y \in \sigma$, $Y_\sigma G_y$ is the retarded Green function for L_g with source δ_y (see [6] for its existence). Thus if f is smooth, then we have the representation formulas $(\gamma_s f)(x) = \int_{\sigma_s} f(y) (Y_{\sigma_s} G_y)(x) \lambda_s(y) d\mu_g(y)$ and $(\Gamma_s f)(x) = \int_{\{y \succ \sigma_s\}} f(y) (Y_{\sigma_s} G_y)(x) d\text{vol}_g(y)$.

Proposition 3.1 *Let (\mathcal{M}, g) be a n -dimensional Lorentzian manifold. Assume that there exist a temporal function τ and a radial function ρ which satisfy (49)–(54). Let $(\sigma_s)_{s \in \mathbb{R}}$ be a $\underline{\sigma}$ -family of admissible Cauchy hypersurfaces s.t. $\lim_{s \rightarrow +\infty} (\inf_{x \in \sigma_s} \tau(x)) = +\infty$. Let $f \in L^1_{loc}(\mathbb{R}, L^2_\ell(\Sigma_\tau))$ s.t. $[s \mapsto \|\lambda_s f|_{\sigma_s}\|_{L^2}] \in L^1_{loc}(\mathbb{R})$.*

Then for any $u \in \mathcal{F}^1_{loc}(\Sigma_\tau)$ s.t. $L_g u = f$, we have, for any $s \in \mathbb{R}$,

$$u = \Theta_{\sigma_s} u + \Gamma_s f \quad \text{on } \{x \succ \sigma_s\}. \quad (66)$$

Moreover

$$\Gamma_s f = \int_s^\infty (\gamma_{s_1} f) ds_1. \quad (67)$$

Remark — The integral in the r.h.s. of (67) makes sense as a distribution on \mathcal{M} since, for any $\varphi \in \mathcal{C}_c^\infty(\mathcal{M})$, we can set $\langle \int_s^\infty (\gamma_{s_1} f) ds_1, \varphi \rangle = \int_s^\infty \langle \gamma_{s_1} f, \varphi \rangle ds_1 = \int_s^{\bar{s}} \langle \gamma_{s_1} f, \varphi \rangle ds_1$, where \bar{s} is s.t. $\text{supp} \varphi \subset \{\sigma_s \prec x \prec \sigma_{\bar{s}}\}$ (\bar{s} exists because $\lim_{s \rightarrow +\infty} (\inf_{x \in \sigma_s} \tau(x)) = +\infty$).

Proof — The proof of (66) is easy: since σ_s is admissible, there exists some $t \in \mathbb{R}$ s.t. $\Sigma_t \prec \sigma_s$ and thus $[\Gamma_s f]_{\Sigma_t} = 0$. Using arguments similar to the ones used in the proofs of Theorem 3.1 or Lemma 3.1, one can deduce that $E_{\Gamma_s f}(\sigma_s) = 0$, i.e. $[\Gamma_s f]_{\sigma_s} = 0$. Hence the Cauchy data on σ_s of both sides of (66) coincide. Since these both sides are also solution of the equation $L_g \varphi = f$ on $\{x \succ \sigma_s\}$, (66) follows by uniqueness of the solution.

To prove (67), fix $s \in \mathbb{R}$ and set $v := \int_s^\infty (\gamma_{s_1} f) ds_1$. We take any $\varphi \in \mathcal{C}_c^\infty(\mathcal{M})$ and compute

$$\int_{\mathcal{M}} (L_g v) \varphi d\text{vol}_g = \int_{\mathcal{M}} v L_g \varphi d\text{vol}_g = \int_{\mathcal{M}} \left(\int_s^\infty (\gamma_{s_1} f) ds_1 \right) L_g \varphi d\text{vol}_g.$$

By Fubini's theorem

$$\int_{\mathcal{M}} (L_g v) \varphi d\text{vol}_g = \int_s^\infty ds_1 \int_{\mathcal{M}} (\gamma_{s_1} f) L_g \varphi d\text{vol}_g = \int_s^\infty ds_1 \int_{x \succ \sigma_{s_1}} (\gamma_{s_1} f) L_g \varphi d\text{vol}_g.$$

Using the identity $\psi L_g \varphi - \varphi L_g \psi = \psi \square_g \varphi - \varphi \square_g \psi = \nabla_\mu (g^{\mu\nu} (\psi \partial_\nu \varphi - \varphi \partial_\nu \psi))$ for $\psi = \gamma_{s_1} f$ and Stokes' theorem we find (taking into account the fact that $\partial\{x \succ \sigma_{s_1}\} = -\sigma_{s_1}$)

$$\begin{aligned} \int_{\mathcal{M}} (L_g v) \varphi d\text{vol}_g &= \int_s^\infty ds_1 \int_{x \succ \sigma_{s_1}} \varphi L_g (\gamma_{s_1} f) d\text{vol}_g \\ &\quad - \int_s^\infty ds_1 \int_{\sigma_{s_1}} \langle N, (\gamma_{s_1} f) \nabla \varphi - \varphi \nabla (\gamma_{s_1} f) \rangle_g d\mu_g \\ &= 0 + \int_s^\infty ds_1 \int_{\sigma_{s_1}} \varphi \langle N, \nabla (\gamma_{s_1} f) \rangle_g d\mu_g = \int_s^\infty ds_1 \int_{\sigma_{s_1}} \varphi \lambda_{s_1} f d\mu_g. \end{aligned}$$

Hence using the definition of λ_s and viewing $d\mu_g$ as a n -form, we deduce that

$$\begin{aligned} \int_{\mathcal{M}} (L_g v) \varphi d\text{vol}_g &= \int_s^\infty ds_1 \int_{\underline{\sigma}} \langle N(F_{s_1}), \frac{\partial F}{\partial s}(s_1, \cdot) \rangle_g F_{s_1}^* (\varphi f d\mu_g) \\ &= \int_s^\infty \int_{\underline{\sigma}} F^* (\varphi f \langle N, \cdot \rangle_g \wedge d\mu_g) \end{aligned}$$

But since on σ_{s_1} , $\langle N, \cdot \rangle_g \wedge d\mu_g = d\text{vol}_g$ (again viewing $d\text{vol}_g$ as a $(n+1)$ -form),

$$\int_{\mathcal{M}} (L_g v) \varphi d\text{vol}_g = \int_s^\infty \int_{\underline{\sigma}} F^* (\varphi f d\text{vol}_g) = \int_{x \succ \sigma_s} \varphi f d\text{vol}_g.$$

which proves $L_g v = f Y_{\sigma_s}$ in the distribution sense. Since we have obviously $v = 0$, for $\{x \prec \sigma_s\}$, we deduce $v = \Gamma_s f$ by uniqueness. Hence (67) follows. \square

3.3 Formulation of the dynamics

We show here a result analogous to Theorem 24 for the nonlinear cubic Klein–Gordon equation

$$\square_g u + u^3 = 0, \tag{68}$$

on a 4-dimensional space-time \mathcal{M} satisfying the hypotheses of Theorem 3.1, involving a smooth family of admissible Cauchy hypersurfaces $(\sigma_s)_{s \in \mathbb{R}}$. We need technical assumptions on $(\sigma_s)_{s \in \mathbb{R}}$, namely:

$$\exists C_1 > 0, \quad \forall s \in \mathbb{R}, \forall x \in \sigma_s, \quad |\lambda_s(x)| \leq C_1 \tag{69}$$

and

$$\exists C_2 > 0, \quad \forall s \in \mathbb{R}, \forall u \in H^1(\sigma_s), \quad \|u\|_{L^6(\sigma_s)} \leq C_2 \|\nabla u\|_{L^2}. \tag{70}$$

Note that (70) is the assumption that the Sobolev embedding $H_0^1(\mathbb{R}^3) \subset L^6(\mathbb{R}^3)$ can be extended on each 3-dimensional manifold σ_s uniformly in s . This is true if e.g. the Ricci curvature of all σ_s is uniformly bounded from below and the volumes of all unit balls in σ_s are uniformly bounded from below (see [22]).

Theorem 3.2 *Let (\mathcal{M}, g) be a 4-dimensional pseudo-Riemannian manifold and $\tau, \rho \in \mathcal{C}^\infty(\mathcal{M})$ satisfying (49)–(54). Let $(\sigma_s)_{s \in \mathbb{R}}$ be a $\underline{\sigma}$ -family of admissible Cauchy hypersurfaces which satisfies (69) and (70) and s.t. $\sup_s C(\sigma_s) < +\infty$. Consider the non autonomous vector field $V : \mathbb{R} \times \mathcal{E}_0^1(\Sigma_\tau) \rightarrow \mathcal{E}_0^1(\Sigma_\tau)$ defined by $V(s, \varphi) := \Phi_{\sigma_s}(0, \lambda_s \varphi^3|_{\sigma_s})$.*

Let $I = [t_1, t_2]$ and J be intervals of \mathbb{R} s.t. $\Sigma_{t_1} \prec \sigma_s \prec \Sigma_{t_2}, \forall s \in J$ and $u \in \mathcal{F}_I^1(\Sigma_\tau)$. If u is a solution of (68), then $\Theta_{\sigma_s} u$ is a \mathcal{C}^1 function of $s \in J$ and satisfies:

$$\frac{d(\Theta_{\sigma_s} u)}{ds} + V(s, \Theta_{\sigma_s} u) = 0, \quad \forall s \in J. \quad (71)$$

Proof — First note that V exists and is continuous because of Corollary 3.1 and of (69) and (70), which imply in particular: $\forall \varphi \in \mathcal{E}_0^1(\Sigma_\tau), \forall s \in \mathbb{R}, \lambda_s \varphi^3|_{\sigma_s} \in L^2(\sigma_s)$. Second let $u \in \mathcal{F}_I^1(\Sigma_\tau)$ and assume that u is a solution of (68).

Step 1 — We show that $[s \mapsto \Theta_{\sigma_s} u]$ is continuous, i.e. $\forall s \in J$,

$$\lim_{s' \rightarrow s} \left(\sup_{t \in I} E_{(\Theta_{\sigma_{s'}} u) - (\Theta_{\sigma_s} u)}(\Sigma_t)^{1/2} \right) = 0.$$

Since $(\Theta_{\sigma_{s'}} u) - (\Theta_{\sigma_s} u) \in \mathcal{E}_0^1(\Sigma_t)$, it suffices to prove $\lim_{s' \rightarrow s} E_{(\Theta_{\sigma_{s'}} u) - (\Theta_{\sigma_s} u)}(\sigma_{s'}) = 0$ and to apply Corollary 3.1 with $\sigma_{s'}$. But actually $[\Theta_{\sigma_{s'}} u]_{\sigma_{s'}} = [u]_{\sigma_{s'}}$, so that $E_{(\Theta_{\sigma_{s'}} u) - (\Theta_{\sigma_s} u)}(\sigma_{s'}) = E_{u - (\Theta_{\sigma_s} u)}(\sigma_{s'})$. Now observe that $[u - (\Theta_{\sigma_s} u)]_{\sigma_s} = 0$ or equivalently $E_{u - (\Theta_{\sigma_s} u)}(\sigma_s) = 0$. Thus in particular the result is straightforward in the case where u is smooth with compact spatial support. The general case follows by proving the existence of a sequence of smooth functions with compact spatial support which converges to u in the $\mathcal{F}_I^1(\Sigma_\tau)$ topology. For that purpose first approach $-u^3$ by a sequence of smooth maps with compact spatial support $(f_\varepsilon)_{\varepsilon > 0}$ in $L^1(I, L_\ell^2(\Sigma_\tau))$ and, for some Cauchy hypersurface Σ , approach $[u]_\Sigma$ by a sequence $(v_\varepsilon, w_\varepsilon)_{\varepsilon > 0}$ of smooth maps with compact support in the $H_m^1(\Sigma) \times L^2(\Sigma)$ topology. For any $\varepsilon > 0$ consider the solution u_ε of $L_g u_\varepsilon = f_\varepsilon$, with the Cauchy data $[u_\varepsilon]_\Sigma = (v_\varepsilon, w_\varepsilon)$. Then u_ε is smooth with compact spatial support and converges to u in $\mathcal{F}_I^1(\Sigma_\tau)$, when $\varepsilon \rightarrow 0$, because of (55).

Step 2 — We use the generalized Duhamel formula. First by applying Lemma 3.1 to u and for $\widehat{\Sigma} = \sigma_s$, we deduce that $s \mapsto \|u|_{\sigma_s}\|_{H_0^1}$ is bounded. Hence again because of (69) and (70), $s \mapsto \|\lambda_s u^3|_{\sigma_s}\|_{L^2}$ is bounded. Thus we can apply Proposition 3.1. Then (66) reads

$$u + \Gamma_s(u^3) = \Theta_{\sigma_s} u \quad \text{on } \{x \succ \sigma_s\}.$$

Comparing this identity for two different value s_1, s_2 of s , we get

$$\Theta_{\sigma_{s_2}} u - \Theta_{\sigma_{s_1}} u = \Gamma_{s_2}(u^3) - \Gamma_{s_1}(u^3) \quad \text{on } \{x \succ \sigma_{s_1}\} \cap \{x \succ \sigma_{s_2}\}. \quad (72)$$

However the r.h.s. of (72) can be written by using (67)

$$\Gamma_{s_2}(u^3) - \Gamma_{s_1}(u^3) = - \int_{s_1}^{s_2} (\gamma_s u^3) ds.$$

Moreover, since $(\Theta_{\sigma_s} u)|_{\sigma_s} = u|_{\sigma_s}$,

$$\gamma_s u^3 = \Phi_{\sigma_s}(0, \lambda_s u^3) = V(s, \Theta_{\sigma_s} u) \quad \text{on } \{x \succ \sigma_s\}$$

Hence (72) implies that the following identity holds on $\{x \succ \sigma_{s_1}\} \cap \{x \succ \sigma_{s_2}\}$:

$$\Theta_{\sigma_{s_2}} u - \Theta_{\sigma_{s_1}} u + \int_{s_1}^{s_2} V(s, \Theta_{\sigma_s} u) ds = 0. \quad (73)$$

But since the l.h.s. of (73) is a solution of $L_g \varphi = 0$ on \mathcal{M} , (73) holds actually everywhere on \mathcal{M} , by uniqueness. From (73), the result of the first step and Corollary 3.1 we then deduce easily (71). \square

4 The space of analytic functions over a Banach space

4.1 Analytic functions over a Banach space

Recall that, if \mathbb{X} and \mathbb{Y} are Banach spaces and $r \in (0, +\infty)$, $\mathbb{F}_r(\mathbb{X}, \mathbb{Y})$ is the space of formal series $f = \sum_{p=0}^{\infty} f^{(p)}$ s.t. $[f](r) < +\infty$, where $[f](z)$ is given by (17). Note that $(\mathbb{F}_r(\mathbb{X}, \mathbb{Y}), [\cdot](r))$ is a Banach space.

Beside the definition of $[f]$ given by (17), we also set, for $k \in \mathbb{N}$,

$$[f]^{(k)}(r) := \frac{d^k}{dz^k} [f](z)|_{z=r} \quad \text{and} \quad \mathbb{F}_r^{(k)}(\mathbb{X}, \mathbb{Y}) := \{f \in \mathbb{F}_r(\mathbb{X}, \mathbb{Y}) \mid [f]^{(k)}(r) < +\infty\},$$

so that for instance $[f]^{(1)}(r) = \sum_{p=1}^{\infty} p \|f^{(p)}\|_{\otimes} r^{p-1}$. We set $\mathbb{F}_{\infty}(\mathbb{X}, \mathbb{Y}) := \bigcap_{r>0} \mathbb{F}_r(\mathbb{X}, \mathbb{Y})$ and $\mathbb{F}_{pol}(\mathbb{X}, \mathbb{Y}) := \{f = \sum_{p=0}^N f^{(p)} \mid N \in \mathbb{N}, f^{(p)} \in \mathcal{Q}^p(\mathbb{X}, \mathbb{Y})\}$. Note that we have the dense inclusions

$$\forall r, R \in (0, \infty), \text{ s.t. } r < R, \forall k, \ell \in \mathbb{N} \text{ s.t. } k < \ell, \quad \mathbb{F}_{pol} \subsetneq \mathbb{F}_{\infty} \subsetneq \mathbb{F}_R \subsetneq \mathbb{F}_r^{(\ell)} \subsetneq \mathbb{F}_r^{(k)} \subset \mathbb{F}_r.$$

Indeed if $0 < r < R$ and $k \in \mathbb{N}$, we have: $\forall f \in \mathbb{F}_R$,

$$[f]^{(k)}(r) \leq \Gamma^{(k)}(r, R) [f](R), \quad \text{where } \Gamma^{(k)}(r, R) := \frac{1}{r^k} \sup_{p \geq k} \frac{p!}{(p-k)!} \left(\frac{r}{R}\right)^p < +\infty. \quad (74)$$

In the following we set $\mathbb{Y} = \mathbb{R}$ and:

Definition 4.1 For any $r_0 \in (0, \infty]$ and any $k, \ell \in \mathbb{N}$ a **continuous operator \mathbb{T} from $\mathbb{F}_{(0, r_0)}^{(k)}(\mathbb{X})$ to $\mathbb{F}_{(0, r_0)}^{(\ell)}(\mathbb{X})$** is a family $(\mathbb{T}_r)_{0 < r < r_0}$, s.t., for any $r \in (0, r_0)$, $\mathbb{T}_r : \mathbb{F}_r^{(k)}(\mathbb{X}) \rightarrow \mathbb{F}_r^{(\ell)}(\mathbb{X})$ is a continuous linear operator with norm $\|\mathbb{T}_r\|$ and s.t., $\forall r, r' \in (0, r_0)$, if $r < r'$, then the restriction of \mathbb{T}_r to $\mathbb{F}_{r'}^{(k)}(\mathbb{X})$ coincides with $\mathbb{T}_{r'}$.

For simplicity we systematically denote each operator \mathbb{T}_r by \mathbb{T} in the following..

4.2 Analytic vector fields over \mathbb{X}

Definition 4.2 Elements of $\mathbb{F}_r(\mathbb{X}, \mathbb{X})$ are called **analytic vector fields on \mathbb{X}** . For any $V \in \mathbb{F}_r(\mathbb{X}, \mathbb{X})$, we denote by $V \cdot$ the linear operator acting on $\mathbb{F}_r(\mathbb{X})$ defined by

$$\forall f \in \mathbb{F}_r(\mathbb{X}), \forall \varphi \in B_{\mathbb{X}}(r), \quad (V \cdot f)(\varphi) = \delta f_{\varphi}(V(\varphi)),$$

where, $\forall \varphi \in B_{\mathbb{X}}(r)$, $\forall \psi \in \mathbb{X}$,

$$\delta f_{\varphi}(\psi) := \lim_{\varepsilon \rightarrow 0} \frac{f(\varphi + \varepsilon \psi) - f(\varphi)}{\varepsilon}$$

We then set $[V] \cdot := [V](z) \frac{d}{dz}$, a holomorphic vector field on $B_{\mathbb{C}}(\rho_V)$.

The previous definition was vague concerning the domain and the target of $V \cdot$. These points are made more precise by the following result.

Lemma 4.1 *For any $V \in \mathbb{F}_r(\mathbb{X}, \mathbb{X})$, the operator $V \cdot$ is continuous from $\mathbb{F}_{(0,r)}^{(1)}(\mathbb{X})$ to $\mathbb{F}_{(0,r)}(\mathbb{X})$ and moreover:*

$$\forall \rho \in (0, r), \quad \forall f \in \mathbb{F}_{\rho}^{(1)}(\mathbb{X}), \quad [V \cdot f](\rho) \leq [V](\rho)[f]^{(1)}(\rho) = ([V] \cdot [f])(\rho). \quad (75)$$

Proof — Consider $\rho \in (0, r)$, assume momentarily that $f \in \mathbb{F}_{\text{pol}}(\mathbb{X})$ and write $f(\varphi) = \sum_{p=0}^N f^{(p)}(\varphi^{\otimes p})$. Then, $\forall \varphi \in \mathbb{X}$ such that $\|\varphi\|_{\mathbb{X}} \leq r$ we know that $V(\varphi)$ is well defined and, using everywhere the convention $p' := p - 1$ and, setting $\varphi_1 \otimes \cdots \otimes \varphi_p = \varphi_1 \cdots \varphi_p$ for short,

$$\begin{aligned} (V \cdot f)(\varphi) &= \delta f_{\varphi}(V(\varphi)) = \sum_{p=1}^N p f^{(p)}(V(\varphi) \underbrace{\varphi \cdots \varphi}_{p'}) \\ &= \sum_{p=1}^N \sum_{q=0}^{\infty} p f^{(p)}(V^{(q)}(\underbrace{\varphi \cdots \varphi}_q \underbrace{\varphi \cdots \varphi}_{p'})) = \sum_{m=0}^{\infty} (V \cdot f)^{(m)}(\underbrace{\varphi \cdots \varphi}_m), \end{aligned}$$

where we have set $m = q + p - 1 = q + p'$ and, $\forall \varphi_1, \dots, \varphi_m \in \mathbb{X}$,

$$(V \cdot f)^{(m)}(\varphi_1 \cdots \varphi_m) := \sum_{p=1}^{\sup(N, m+1)} \frac{1}{m!} \sum_{\sigma \in \mathfrak{S}_m} p f^{(p)} \left(V^{(m-p')}(\varphi_{\sigma(1)} \cdots \varphi_{\sigma(m-p')}) \varphi_{\sigma(m-p'+1)} \cdots \varphi_{\sigma(m)} \right).$$

Hence $|(V \cdot f)^{(m)}(\varphi_1 \cdots \varphi_m)|$ is less than or equal to (we set $\|\cdot\| = \|\cdot\|_{\mathbb{X}}$ for shortness):

$$\sum_{p=1}^{\sup(N, m+1)} \frac{1}{m!} \sum_{\sigma \in \mathfrak{S}_m} p \|f^{(p)}\|_{\otimes} \left\| V^{(m-p')}(\varphi_{\sigma(1)} \cdots \varphi_{\sigma(m-p')}) \right\| \|\varphi_{\sigma(m-p'+1)}\| \cdots \|\varphi_{\sigma(m)}\| \quad (76)$$

and since $\left\| V^{(m-p')}(\varphi_{\sigma(1)} \cdots \varphi_{\sigma(m-p')}) \right\| \leq \|V^{(m-p')}\|_{\otimes} \|\varphi_{\sigma(1)}\| \cdots \|\varphi_{\sigma(m-p')}\|$, we deduce from the upper bound (76) that

$$\begin{aligned} |(V \cdot f)^{(m)}(\varphi_1 \cdots \varphi_m)| &\leq \sum_{p=1}^{\sup(N, m+1)} \frac{1}{m!} \sum_{\sigma \in \mathfrak{S}_m} p \|V^{(m-p')}\|_{\otimes} \|f^{(p)}\|_{\otimes} \|\varphi_{\sigma(1)}\| \cdots \|\varphi_{\sigma(m)}\| \\ &= \sum_{p=1}^{\sup(N, m+1)} p \|V^{(m-p')}\|_{\otimes} \|f^{(p)}\|_{\otimes} \|\varphi_1\| \cdots \|\varphi_m\|. \end{aligned}$$

We thus deduce

$$\|(V \cdot f)^{(m)}\|_{\otimes} \leq \sum_{p=1}^{\sup(N, m+1)} p \|V^{(m-p')}\|_{\otimes} \|f^{(p)}\|_{\otimes}. \quad (77)$$

Hence, by letting $q = m - p'$,

$$\begin{aligned} [V \cdot f](r) &= \sum_{m=0}^{\infty} \|(V \cdot f)^{(m)}\|_{\otimes} r^m \leq \sum_{m=0}^{\infty} \sum_{p=1}^{\sup(N, m+1)} p \|V^{(m-p')}\|_{\otimes} \|f^{(p)}\|_{\otimes} r^m \\ &= \sum_{q=0}^{\infty} \sum_{p=1}^N \|V^{(q)}\|_{\otimes} r^q p \|f^{(p)}\|_{\otimes} r^{p-1} = [V](r)[f]^{(1)}(r). \end{aligned}$$

Thus we obtain (75) for $f \in \mathbb{F}_{pol}(\mathbb{X})$. It implies the result by using the density of $\mathbb{F}_{pol}(\mathbb{X})$ in $\mathbb{F}_r^{(1)}(\mathbb{X})$. \square

Note that we can extend (77) *a posteriori* to any $f \in \mathbb{F}_r^{(1)}(\mathbb{X})$ by density as soon as $\sum_{q=0}^{\infty} \|V^{(q)}\| r^q < +\infty$, thanks to (75). It gives us (still with the convention $p' = p - 1$):

$$\|(V \cdot f)^{(m)}\|_{\otimes} \leq \sum_{p=1}^m p \|V^{(m-p')}\|_{\otimes} \|f^{(p)}\|_{\otimes}. \quad (78)$$

This leads us to the following extension of Lemma 4.1.

Lemma 4.2 *Let $k \in \mathbb{N}^*$, $r_0 > 0$ and $V_1, \dots, V_k \in \mathbb{F}_{r_0}(\mathbb{X}, \mathbb{X})$. Then the linear operator $[f \mapsto V_k \cdots V_1 \cdot f]$ is continuous from $\mathbb{F}_{(0, r_0)}^{(k)}(\mathbb{X})$ to $\mathbb{F}_{(0, r_0)}(\mathbb{X})$ and $\forall r \in (0, r_0)$,*

$$\forall f \in \mathbb{F}_r^{(k)}, \quad [V_k \cdots V_1 \cdot f](r) \leq ([V_k] \cdots [V_1] \cdot [f])(r). \quad (79)$$

Proof — For any $a = 1, \dots, k$ we write $V_a = \sum_{p=0}^{\infty} V_a^{(p)}$, where $\forall p \in \mathbb{N}$, $V_a^{(p)} \in \mathcal{Q}^p(\mathbb{X}, \mathbb{X})$. For shortness we set $X_a^{(p)} := \|V_a^{(p)}\|_{\otimes}$, $X_a(z) := \sum_{p \geq 0} X_a^{(p)} z^p$ and $X_a \cdot := X_a(z) \frac{d}{dz}$. We recall that $\forall f \in \mathbb{F}, \forall \varphi \in \mathbb{X}$, $(V_a \cdot f)(\varphi) = \delta f_{\varphi}(V_a(\varphi))$. In the following we assume first that $f \in \mathbb{F}_{pol}$. On the one hand we observe that, $\forall p \in \mathbb{N}$,

$$\|(V_k \cdots V_1 \cdot f)^{(p)}\|_{\otimes} \leq \sum_{p'_k=0}^p \sum_{p'_{k-1}=0}^{p_k} \cdots \sum_{p'_1=0}^{p_2} p_k \cdots p_1 X_k^{(p-p'_k)} X_{k-1}^{(p_k-p'_{k-1})} \cdots X_1^{(p_2-p'_1)} \|f^{(p_1)}\|_{\otimes}, \quad (80)$$

where we systematically denote $p'_a := p_a - 1$. This can be proved by recursion on k , by using (78). On the other hand the coefficients of the decomposition $(X_k \cdots X_1 \cdot [f])(z) = \sum_{p=0}^{\infty} (X_k \cdots X_1 \cdot [f])^{(p)} z^p$ also satisfy similar relations, i.e.

$$(X_k \cdots X_1 \cdot [f])^{(p)} = \sum_{p'_k=0}^p \sum_{p'_{k-1}=0}^{p_k} \cdots \sum_{p'_1=0}^{p_2} p_k \cdots p_1 X_k^{(p-p'_k)} X_{k-1}^{(p_k-p'_{k-1})} \cdots X_1^{(p_2-p'_1)} \|f^{(p_1)}\|_{\otimes},$$

which can also be proved by a recursion based on the identity

$$X_a \cdot \left(\sum_{p=0}^{\infty} A^{(p)} z^p \right) = \sum_{m=0}^{\infty} \left(\sum_{p'=0}^m p X_a^{(m-p')} A^{(p)} \right) z^m.$$

Hence the result follows easily from this identity and (80) holds for $f \in \mathbb{F}_{pol}(\mathbb{X})$. This can hence be extended to all $f \in \mathbb{F}_r^{(k)}(\mathbb{X})$ for $r \in (0, r_0)$ by density. \square

5 The time ordered exponential of operators

In this section we consider a Lebesgue measurable family $(V_t \cdot)_{t \in I}$ of continuous operators $V_t \cdot$ from $\mathbb{F}_{(0, r_0)}^{(1)}(\mathbb{X})$ to $\mathbb{F}_{(0, r_0)}(\mathbb{X})$ and we consider the time ordered exponential

$$U_{t_1}^{t_2} := T \exp \int_{t_1}^{t_2} d\tau (V_\tau \cdot) := \sum_{k=0}^{\infty} \frac{(V \cdot)_{t_1}^{t_2[k]}}{k!}, \quad (81)$$

where $(V \cdot)_{t_1}^{t_2[0]} := 1_{\text{End}(\mathbb{F})}$ and for $k \geq 1$,

$$(V \cdot)_{t_1}^{t_2[k]} := k! \int_{t_1 < \tau_1 < \dots < \tau_k < t_2} (V_{\tau_k} \cdots V_{\tau_1} \cdot) d\tau_1 \cdots d\tau_k, \quad \text{for } t_2 > t_1 \quad (82)$$

and

$$(V \cdot)_{t_1}^{t_2[k]} := (-1)^k k! \int_{t_2 < \tau_k < \dots < \tau_1 < t_1} (V_{\tau_k} \cdots V_{\tau_1} \cdot) d\tau_1 \cdots d\tau_k, \quad \text{for } t_2 < t_1. \quad (83)$$

We remark that, for $t_2 > t_1$,

$$\frac{(V \cdot)_{t_1}^{t_2[k]}}{k!} = \int_{t_1}^{t_2} d\tau_k V_{\tau_k} \cdot \left(\int_{t_1 < \tau_1 < \dots < \tau_{k-1} < \tau_k} (V_{\tau_{k-1}} \cdots V_{\tau_1} \cdot) d\tau_1 \cdots d\tau_{k-1} \right) = \int_{t_1}^{t_2} d\tau V_\tau \cdot \frac{(V \cdot)_{t_1}^{\tau[k-1]}}{(k-1)!}.$$

Hence

$$U_{t_1}^{t_2} = 1_{\text{End}(\mathbb{F})} + \int_{t_1}^{t_2} d\tau V_\tau \cdot U_{t_1}^\tau. \quad (84)$$

A similar reasoning shows that (84) holds also for $t_2 < t_1$. As a consequence

$$U_{t_1}^{t_2+h} - U_{t_1}^{t_2} = \int_{t_1}^{t_2+h} d\tau V_\tau \cdot U_{t_1}^\tau - \int_{t_1}^{t_2} d\tau V_\tau \cdot U_{t_1}^\tau = \int_{t_2}^{t_2+h} d\tau V_\tau \cdot U_{t_1}^\tau. \quad (85)$$

5.1 Existence of $U_{t_1}^{t_2}$

In the following, for any vector field X on $B_{\mathbb{C}}(r)$, we denote by $(t, z) \mapsto e^{-tX}(z)$ the map which is equal to the solution γ of

$$\begin{cases} \frac{\partial \gamma}{\partial t}(t, z) &= -X(\gamma(t, z)) \\ \gamma(0, z) &= z. \end{cases}$$

Theorem 5.1 *Let $r_0 \in (0, +\infty]$ and $I \subset \mathbb{R}$ be an interval. Let $(V_t)_{t \in I}$ be a normal family of analytic vector fields in $\mathbb{F}_{r_0}(\mathbb{X}, \mathbb{X})$ and let $X = \sum_{k=0}^{\infty} X_k z^k \in \mathbb{F}_{r_0}(\mathbb{R})$ s.t. $\forall t \in I, \forall p \in \mathbb{N}, 0 \leq [V_t^{(p)}] \leq X_p$. Assume that:*

$$\forall r \in (0, r_0), \forall f \in \mathbb{F}_r^{(1)}(\mathbb{X}), \quad [I \ni t \mapsto V_t \cdot f \in \mathbb{F}_r(\mathbb{X})] \text{ is measurable.} \quad (86)$$

Let $R \in (0, r_0)$. Then $\forall t_1, t_2 \in I$ s.t. $e^{-|t_2-t_1|X}(R)$ exists and is positive, the operator $U_{t_1}^{t_2} := T \exp\left(\int_{t_1}^{t_2} d\tau V_\tau \cdot\right)$ defined by (81) is a bounded operator from $\mathbb{F}_R(\mathbb{X})$ to $\mathbb{F}_{e^{-|t_2-t_1|X}(R)}(\mathbb{X})$ with a norm less than 1 i.e.

$$\forall t \in [t_1, t_2], \forall f \in \mathbb{F}_R(\mathbb{X}), \quad [U_{t_1}^{t_2} f](e^{-|t_2-t_1|X}(R)) \leq [f](R). \quad (87)$$

Moreover for any \bar{R} s.t. $R < \bar{R} < r_0$ and $f \in \mathbb{F}_{\bar{R}}(\mathbb{X})$, the map $t \mapsto U_{t_1}^t f$ is locally Lipschitz continuous from $[t_1, t_2]$ to $\mathbb{F}_{e^{-|t_2-t_1|X}(R)}(\mathbb{X})$.

Proof of theorem 5.1 — W.l.g. we assume throughout the proof that $t_1 = 0 < T = t_2$ and study U_0^t for $0 < t \leq T$. The proof is divided in several steps which follow.

Step 1 — For $r \in (0, r_0)$, $k \in \mathbb{N}$, $f \in \mathbb{F}(\mathbb{X})$ and $t \in I$ we estimate the norm in $\mathbb{F}_r(\mathbb{X})$ of $(V \cdot)_0^{t[k]} f$. For $t \in [0, T]$, we start from Expression (82) for $(V \cdot)_0^{t[k]} f$ and we use Lemma 4.2 with $V_a \cdot = V_{\tau_a} \cdot$. This gives us

$$\begin{aligned} [(V \cdot)_0^{t[k]} f](r) &\leq k! \int_{0 < \tau_1 < \dots < \tau_k < t} [V_{\tau_k} \dots V_{\tau_1} \cdot f](r) d\tau_1 \dots d\tau_k \\ &\leq k! \int_{0 < \tau_1 < \dots < \tau_k < t} ([V_{\tau_k}] \dots [V_{\tau_1}] \cdot [f])(r) d\tau_1 \dots d\tau_k \\ &\leq k! \int_{0 < \tau_1 < \dots < \tau_k < t} (X \cdot)^k [f](r) d\tau_1 \dots d\tau_k, \end{aligned}$$

which implies, by using $k! \int_{0 < \tau_1 < \dots < \tau_k < t} d\tau_1 \dots d\tau_k = t^k$:

$$[(V \cdot)_0^{t[k]} f](r) \leq t^k (X \cdot)^k [f](r). \quad (88)$$

Hence we see how to derive a sufficient condition for the series $U_0^t f = \sum_{k=0}^{\infty} \frac{1}{k!} (V \cdot)_0^{t[k]} f$ to be convergent in some space \mathbb{F}_r : it suffices to find some r which satisfies

$$\sum_{k=0}^{\infty} \frac{t^k}{k!} (X \cdot)^k [f](r) < +\infty. \quad (89)$$

Then this implies by (88) that $\sum_{k=0}^{\infty} \frac{1}{k!} [(V \cdot)_0^{t[k]} f](r) < +\infty$ and hence the existence of $U_0^t f$.

Step 2 — We show that, if $R \in (0, r_0)$, $T > 0$ and $e^{-TX}(R) > 0$, condition (89) is satisfied with $t = T$ and $r = e^{-TX}(R)$. Actually we will show that

$$\forall f \in \mathbb{F}(\mathbb{X}), \quad \sum_{k=0}^{\infty} \frac{T^k}{k!} ((X \cdot)^k [f]) (e^{-TX}(R)) = [f](R). \quad (90)$$

For that purpose we use the following lemma, the proof of which is given below. In the following, for $t, r > 0$, we set $\bar{B}_{\mathbb{C}}(r) := \{z \in \mathbb{C} \mid |z| \leq r\}$.

Lemma 5.1 *Let $X : B_{\mathbb{C}}(r_0) \mapsto \mathbb{C}$ be an holomorphic vector field different from 0. Assume that*

$$X(z) = \sum_{k=0}^{\infty} X_k z^k, \quad \text{where } X_k \geq 0, \forall k \in \mathbb{N}.$$

Let $\rho \in (0, r_0)$ and $T > 0$ such that $e^{TX}(\rho)$ exists. Then the flow map

$$\begin{aligned} \overline{B}_{\mathbb{C}}(T) \times \overline{B}_{\mathbb{C}}(\rho) &\longrightarrow \mathbb{C} \\ (\tau, z) &\longmapsto e^{\tau X}(z) \end{aligned}$$

is well defined and holomorphic and in particular

$$\forall (\tau, z) \in \overline{B}_{\mathbb{C}}(T) \times \overline{B}_{\mathbb{C}}(\rho), \quad |e^{\tau X}(z)| \leq e^{|\tau|X}(|z|) \leq e^{TX}(\rho). \quad (91)$$

Consider any $R \in (0, r_0)$, $0 < t \leq T$ s.t. $e^{-tX}(R) > 0$: then $e^{tX}(e^{-tX}(R))$ exists since it is nothing but R . Hence we can apply Lemma 5.1 with $\rho = e^{-TX}(R)$. It implies in particular that, for any holomorphic function H on $\overline{B}_{\mathbb{C}}(R) = \overline{B}_{\mathbb{C}}(e^{TX}(\rho))$, the map

$$\begin{aligned} \overline{B}_{\mathbb{C}}(T) \times \overline{B}_{\mathbb{C}}(e^{-TX}(R)) &\longrightarrow \mathbb{C} \\ (\tau, z) &\longmapsto H(e^{\tau X}(z)) \end{aligned}$$

is well defined and is analytic. Hence the following expansion holds:

$$\forall (\tau, z) \in \overline{B}_{\mathbb{C}}(T) \times \overline{B}_{\mathbb{C}}(e^{-TX}(R)), \quad H(e^{\tau X}(z)) = \sum_{k=0}^{\infty} \frac{d^k H(e^{sX}(z))}{(ds)^k} \Big|_{s=0} \frac{\tau^k}{k!}, \quad (92)$$

the series on the r.h.s. being absolutely convergent for any $\tau \in \overline{B}_{\mathbb{C}}(T)$. However because of the identity $(\frac{d}{ds})^k [H(e^{sX}(z))] = ((X \cdot)^k H)(e^{sX}(z))$, which can be proved by recursion over k , we deduce from (92) that

$$\forall (\tau, z) \in \overline{B}_{\mathbb{C}}(T) \times \overline{B}_{\mathbb{C}}(e^{-TX}(R)), \quad H(e^{\tau X}(z)) = \sum_{k=0}^{\infty} ((X \cdot)^k H)(z) \frac{\tau^k}{k!}.$$

By specializing this relation to $(\tau, z) = (T, e^{-TX}(R))$ we deduce that the power series $\sum_{k=0}^{\infty} \frac{T^k}{k!} ((X \cdot)^k H)(e^{-TX}(R))$ is absolutely convergent and satisfies the identity

$$H(R) = \sum_{k=0}^{\infty} \frac{T^k}{k!} ((X \cdot)^k H)(e^{-TX}(R)). \quad (93)$$

Hence by using (93) with $H(z) = [f](z)$ we obtain (90). This shows that the series $T \exp\left(\int_0^T d\tau V_{\tau} \cdot\right) f$ converges in $\mathbb{F}_{e^{-TX}(R)}(\mathbb{X})$. Moreover we deduce using (88) and (90) the following estimate:

$$\left[T \exp\left(\int_0^T d\tau V_{\tau} \cdot\right) f \right] (e^{-TX}(R)) \leq \sum_{k=0}^{\infty} \frac{1}{k!} \left[(V \cdot)_0^{T[k]} f \right] (e^{-TX}(R)) \leq [f](R). \quad (94)$$

Lastly we remark that hypothesis $e^{-TX}(R) > 0$ obviously implies $e^{-tX}(R) > 0, \forall t \in [0, T]$ so that Conclusion (94) holds also if we replace T by $t \in [0, T]$. This implies (87).

Step 3 — Let us prove the local Lipschitz continuity of $t \mapsto U_0^t f$, for $f \in \mathbb{F}_{\overline{R}}(\mathbb{X})$, where $R < \overline{R} < r_0$. Let $t \in [0, T]$ and $h \in \mathbb{R}$ s.t. $t + h \in [0, T]$. Then it follows from (85) and (75) that

$$\begin{aligned} [(U_0^{t+h} - U_0^t)f](e^{-TX}(R)) &\leq \int_t^{t+h} d\tau [V_\tau \cdot U_0^\tau f](e^{-TX}(R)) \\ &\leq |h|X(e^{-TX}(R)) \sup_{t < \tau < t+h} [U_0^\tau f]^{(1)}(e^{-TX}(R)). \end{aligned}$$

However by observing that $e^{-tX}(R) < e^{-tX}(\overline{R})$ because $R < \overline{R}$ we deduce from (74) that $[g]^{(1)}(e^{-TX}(R)) \leq \Gamma^{(1)}(R, \overline{R}) [g](e^{-TX}(\overline{R}))$, $\forall g \in \mathbb{F}_{e^{-TX}(\overline{R})}(\mathbb{X})$. Applying this for $g = U_0^\tau f$,

$$[(U_0^{t+h} - U_0^t)f](e^{-TX}(R)) \leq |h|X(e^{-TX}(R)) \Gamma^{(1)}(R, \overline{R}) \sup_{t < \tau < t+h} [U_0^\tau f](e^{-TX}(\overline{R}))$$

and by using (87) with \overline{R} instead of R :

$$[(U_0^{t+h} - U_0^t)f](e^{-TX}(R)) \leq |h|X(e^{-TX}(R)) \Gamma^{(1)}(R, \overline{R}) [f](\overline{R}).$$

□

Proof of lemma 5.1 — We first show that $(\tau, z) \mapsto e^{\tau X}(z)$ is defined and satisfies (91) over $B_{\mathbb{C}}(T) \times B_{\mathbb{C}}(\rho)$. Fix some $z \in B_{\mathbb{C}}(\rho)$ and $\tau \in B_{\mathbb{C}}(T)$. Then $\exists \varepsilon_0 > 0$ s.t. $\forall \varepsilon \in (0, \varepsilon_0]$,

$$|z| \leq \rho - \varepsilon \quad \text{and} \quad |\tau| \leq T_\varepsilon := \frac{T}{1 + \varepsilon}.$$

We also let $\lambda \in S^1 \subset \mathbb{C}$ such that $\tau = |\tau|\lambda$, where $0 < |\tau| \leq T_\varepsilon$. We introduce the notations:

$$\begin{cases} f_\varepsilon(t) &:= e^{t(1+\varepsilon)X}(|z| + \varepsilon) & \forall t \in [0, T_\varepsilon] \\ \gamma(t) &:= e^{t\lambda X}(z) & \forall t \in [0, \bar{t}] \\ g(t) &:= |\gamma(t)| & \forall t \in [0, \bar{t}] \end{cases}$$

where \bar{t} is the positive maximal existence time for γ . Note that f_ε is defined on $[0, T_\varepsilon]$ because of the assumption that $e^{TX}(R)$ exists. Our first task is to show that the set:

$$A_\varepsilon := \{t \in [0, T_\varepsilon] \cap [0, \bar{t}] \mid g(t) - f_\varepsilon(t) \geq 0\}$$

is actually empty. Let us prove it by contradiction and assume that $A_\varepsilon \neq \emptyset$. Then there exists $t_0 := \inf A_\varepsilon$. Note that $g(0) - f_\varepsilon(0) = -\varepsilon < 0$, hence we deduce from the continuity of $g - f_\varepsilon$ that $t_0 \neq 0$ and $g(t_0) = f_\varepsilon(t_0)$. Moreover since $f_\varepsilon(0) = \varepsilon$ and f_ε is increasing because $X(r) > 0$ for $r > 0$ we certainly have $g(t_0) = f_\varepsilon(t_0) > 0$. We now observe that

$$\forall z \in \mathbb{C}^*, \quad \frac{\langle \lambda X(z), z \rangle}{|z|} = \left\langle \lambda \sum_{k=0}^{\infty} X_k z^k, \frac{z}{|z|} \right\rangle \leq \sum_{k=0}^{\infty} X_k |z|^k = X(|z|).$$

Hence for all $t \geq 0$ s.t. $g(t) \neq 0$,

$$g'(t) = \frac{\langle \lambda X(\gamma(t)), \gamma(t) \rangle}{|\gamma(t)|} \leq X(|\gamma(t)|) = X(g(t))$$

and hence in particular, since $g(t_0) \neq 0$,

$$g'(t_0) \leq X(g(t_0)) = X(f_\varepsilon(t_0)) = \frac{f'_\varepsilon(t_0)}{1 + \varepsilon} < f'_\varepsilon(t_0).$$

Thus since $f'_\varepsilon - g'$ is continuous $\exists t_1 \in (0, t_0)$ s.t. $\forall t \in [t_1, t_0]$, $f'_\varepsilon(t) - g'(t) \geq 0$. Integrating this inequality over $[t_1, t_0]$ we obtain

$$g(t_1) - f_\varepsilon(t_1) = (f_\varepsilon(t_0) - g(t_0)) - (f_\varepsilon(t_1) - g(t_1)) = \int_{t_1}^{t_0} (f'_\varepsilon(t) - g'(t)) dt \geq 0,$$

i.e. $t_1 \in A_\varepsilon$, a contradiction.

Hence $A_\varepsilon = \emptyset$. Note that this implies automatically that $\bar{t} > T_\varepsilon$. Indeed if we had $\bar{t} \leq T_\varepsilon$ this would imply that g is not bounded in $[0, \bar{t}] \subset [0, T_\varepsilon]$, but since f_ε is bounded on $[0, T_\varepsilon]$ we could then find some time $t \in [0, \bar{t}]$ s.t. $g(t) \geq f_\varepsilon(t)$, which would contradict the fact that $A_\varepsilon = \emptyset$. Thus we deduce that $\forall t \in [0, T_\varepsilon]$, $g(t) < f_\varepsilon(t)$, i.e.

$$\forall t \in [0, T_\varepsilon], \quad |e^{\lambda t X}(z)| < e^{(1+\varepsilon)tX}(|z| + \varepsilon).$$

In other words for all $\tau = \lambda t \in B_{\mathbb{C}}(T)$ and all $z \in B_{\mathbb{C}}(\rho)$ we found that $\forall \varepsilon \in (0, \varepsilon_0]$, $|e^{\tau X}(z)| \leq e^{(1+\varepsilon)|\tau|X}(|z| + \varepsilon)$. Letting ε goes to 0, we deduce the estimate (91) for $(\tau, z) \in B_{\mathbb{C}}(T) \times B_{\mathbb{C}}(\rho)$. Lastly this estimate forbids the flow to blow up on $\overline{B_{\mathbb{C}}(T)} \times \overline{B_{\mathbb{C}}(\rho)}$. Hence the result and (91) can be extended to this domain by continuity. \square

6 Proof of the Main Theorem

We first prove the following strengthening of Theorem 5.1 (with stronger hypotheses).

Theorem 6.1 *Let $r_0 \in (0, +\infty]$ and $I \subset \mathbb{R}$ be an interval. Let $(V_t)_{t \in I}$ be a normal family of analytic vector fields in $\mathbb{F}_{r_0}(\mathbb{X}, \mathbb{X})$ and let $X = \sum_{k=0}^{\infty} X_k z^k \in \mathbb{F}_{r_0}(\mathbb{R})$ s.t. $\forall t \in I$, $\forall p \in \mathbb{N}$, $0 \leq [V_t^{(p)}] \leq X_p$. Assume that:*

$$I \times B_{\mathbb{X}}(r_0) \ni (t, \varphi) \longmapsto V_t(\varphi) \in \mathbb{X} \text{ is continuous.} \quad (95)$$

Let $R, \bar{R} \in \mathbb{R}$ s.t. $0 < R < \bar{R} < r_0$. Let $f \in \mathbb{F}_{\bar{R}}(\mathbb{X})$. Let $t_1, t_2 \in I$ s.t. $e^{-|t_2 - t_1|X}(R) > 0$ and let $\varphi \in \mathcal{C}^1([t_1, t_2], \mathbb{X})$ s.t. $\|\varphi(t)\|_{\mathbb{X}} \leq e^{-|t - t_1|X}(R)$, $\forall t \in [t_1, t_2]$. Then the map

$$\begin{aligned} [t_1, t_2] &\longrightarrow \mathbb{X} \\ t &\longmapsto (U_{t_1}^t f)(\varphi(t)) \end{aligned}$$

is \mathcal{C}^1 and satisfies

$$\frac{d}{dt} ((U_{t_1}^t f)(\varphi(t))) = (V_t \cdot U_{t_1}^t f)(\varphi(t)) + \delta(U_{t_1}^t f)_{\varphi(t)} \left(\frac{d\varphi(t)}{dt} \right). \quad (96)$$

Proof — W.l.g. we assume $t_1 = 0 < T = t_2$. Let $f \in \mathbb{F}_{\overline{R}}(\mathbb{X})$ and, for $t \in [0, T]$, set $f_t := U_0^t f$. By Theorem 5.1 we know that $f_t \in \mathbb{F}_{e^{-tX}(\overline{R})}(\mathbb{X})$, $\forall t \in [0, T]$. We first show that, $\forall t \in [0, T]$, $(\tau, \varphi) \mapsto (V_\tau \cdot f_\tau)(\varphi)$ is continuous on $[0, t] \times B_{\mathbb{X}}(0, e^{-tX}(R))$. For that purpose, for $\tau, \tau + \sigma \in [0, t]$ and $\varphi, \psi \in B_{\mathbb{X}}(0, e^{-tX}(R))$ we evaluate the difference $(V_{\tau+\sigma} \cdot f_{\tau+\sigma})(\psi) - (V_\tau \cdot f_\tau)(\varphi)$. We split this quantity as the sum of three terms:

$$(V_{\tau+\sigma} \cdot f_{\tau+\sigma})(\psi) - (V_\tau \cdot f_\tau)(\varphi) = \delta(f_{\tau+\sigma})_\psi(V_{\tau+\sigma}(\psi)) - \delta(f_\tau)_\varphi(V_\tau(\varphi)) = \Delta_1 + \Delta_2 + \Delta_3,$$

where

$$\begin{aligned} \Delta_1 &:= \delta(f_{\tau+\sigma})_\psi(V_{\tau+\sigma}(\psi) - V_\tau(\varphi)) \\ \Delta_2 &:= (\delta(f_{\tau+\sigma})_\psi - \delta(f_{\tau+\sigma})_\varphi)(V_\tau(\varphi)) \\ \Delta_3 &:= \delta(f_{\tau+\sigma} - f_\tau)_\varphi(V_\tau(\varphi)). \end{aligned}$$

To evaluate Δ_1 and Δ_3 we will use the following inequality (for all $r > 0$):

$$\forall g \in \mathbb{F}_r^{(1)}(\mathbb{X}), \forall \varphi \in B_{\mathbb{X}}(r), \forall Z \in \mathbb{X}, \quad |\delta g_\varphi(Z)| \leq [Z \cdot g](\|\varphi\|_{\mathbb{X}}) = \|Z\|_{\mathbb{X}} [g]^{(1)}(\|\varphi\|_{\mathbb{X}}), \quad (97)$$

which follows by applying Lemma 4.1, (75) with V being the constant vector field $[\varphi \mapsto Z]$.

We note that (97), $\|\psi\|_{\mathbb{X}} < e^{-tX}(R)$ and Inequality (74) imply

$$\begin{aligned} |\Delta_1| &\leq \|V_{\tau+\sigma}(\psi) - V_\tau(\varphi)\|_{\mathbb{X}} [f_{\tau+\sigma}]^{(1)}(\|\psi\|_{\mathbb{X}}) \\ &\leq \|V_{\tau+\sigma}(\psi) - V_\tau(\varphi)\|_{\mathbb{X}} \Gamma^{(1)}(e^{-tX}(R), e^{-tX}(\overline{R})) [f_{\tau+\sigma}](e^{-tX}(\overline{R})) \end{aligned}$$

and hence Δ_1 converges to 0 as $\sigma \rightarrow 0$ and $\|\psi - \varphi\|_{\mathbb{X}} \rightarrow 0$ because of (95). We decompose and split Δ_2 :

$$\begin{aligned} \Delta_2 &:= \sum_{p=0}^{\infty} p f_{\tau+\sigma}^{(p)}(V_\tau(\varphi) \underbrace{\psi \cdots \psi}_{p-1}) - p f_{\tau+\sigma}^{(p)}(V_\tau(\varphi) \underbrace{\varphi \cdots \varphi}_{p-1}) \\ &= \sum_{p=0}^{\infty} p \sum_{j=1}^{p-1} f_{\tau+\sigma}^{(p)}(V_\tau(\varphi) \underbrace{\psi \cdots \psi}_{j-1} (\psi - \varphi) \underbrace{\varphi \cdots \varphi}_{p-1-j}) \end{aligned}$$

We deduce that, by setting $M := \sup(\|\varphi\|_{\mathbb{X}}, \|\psi\|_{\mathbb{X}})$,

$$|\Delta_2| \leq \|V_\tau(\varphi)\|_{\mathbb{X}} \|\psi - \varphi\|_{\mathbb{X}} \sum_{p=0}^{\infty} p(p-1) \|f_{\tau+\sigma}^{(p)}\|_{\otimes} M^{p-2} = \|V_\tau(\varphi)\|_{\mathbb{X}} \|\psi - \varphi\|_{\mathbb{X}} [f_{\tau+\sigma}]^{(2)}(M).$$

Hence by using $M \leq e^{-tX}(R) < e^{-tX}((R + \overline{R})/2)$ and Inequality (74), we deduce that Δ_2 tends to 0 when $\|\psi - \varphi\|_{\mathbb{X}} \rightarrow 0$. Lastly using again (97) we have

$$|\Delta_3| \leq \|V_\tau(\varphi)\|_{\mathbb{X}} [f_{\tau+\sigma} - f_\tau]^{(1)}(\|\varphi\|_{\mathbb{X}}),$$

which implies also that Δ_3 tends to 0 when $\sigma \rightarrow 0$ by applying Theorem 5.1 with $(R + \overline{R})/2$ in place of R (since $(R + \overline{R})/2 < \overline{R}$ and $f \in \mathbb{F}_{\overline{R}}$ the map $\tau \mapsto f_\tau$ is continuous from $[0, t]$ to $\mathbb{F}_{e^{-tX}(\frac{R+\overline{R}}{2})}(\mathbb{X})$ and hence to $\mathbb{F}_{e^{-tX}(R)}^{(1)}(\mathbb{X})$ by Inequality (74)).

Hence we conclude that $(V_{\tau+\sigma} \cdot f_{\tau+\sigma})(\psi) - (V_\tau \cdot f_\tau)(\varphi)$ converges to 0 when $\sigma \rightarrow 0$ and $\|\psi - \varphi\| \rightarrow 0$, which proves the continuity of $(\tau, \varphi) \mapsto (V_\tau \cdot f_\tau)(\varphi)$.

An easy consequence is that the r.h.s. of (96) is continuous. Thus it suffices to prove (96) in order to conclude. Let $h \neq 0$, then using (85):

$$\begin{aligned} \frac{1}{h} [f_{t+h}(\varphi(t+h)) - f_t(\varphi(t))] &= \frac{1}{h} [f_{t+h}(\varphi(t+h)) - f_t(\varphi(t+h))] + \frac{1}{h} [f_t(\varphi(t+h)) - f_t(\varphi(t))] \\ &= \frac{1}{h} \int_t^{t+h} d\tau (V_\tau \cdot f_\tau)(\varphi(t+h)) + \frac{1}{h} (f_t(\varphi(t+h)) - f_t(\varphi(t))) \end{aligned} \tag{98}$$

When $h \rightarrow 0$ the first term in the r.h.s. of (98) converges to $(V_t \cdot f_t)(\varphi(t))$ because of the continuity of $(\tau, \varphi) \mapsto (V_\tau \cdot f_\tau)(\varphi)$. The second term in the r.h.s. of (98) converges to $\delta(f_t)_{\varphi(t)} \left(\frac{d\varphi(t)}{dt} \right)$ because of (30). Hence the r.h.s. of (98) converges to $(V_t \cdot f_t)(\varphi(t)) + \delta(f_t)_{\varphi(t)} \left(\frac{d\varphi(t)}{dt} \right)$ when $h \rightarrow 0$, which proves (96). \square

Proof of the Theorem 0.2 — On a flat space-time with a general real analytic nonlinearity we first use Proposition 2.1 which provides us with a normal family of analytic vector fields $(V_t)_{t \in I}$ satisfying (95) and using Theorem 2.1 we obtain a \mathcal{C}^1 map $\varphi(t) = \Theta_t u$ which satisfies (24). We can thus apply Theorem 6.1 to these data and deduce:

$$\frac{d}{dt} ((U_{t_1}^t f)(\Theta_t u)) = (V_t \cdot U_{t_1}^t f)(\Theta_t u) + \delta(U_{t_1}^t f)_{\varphi(t)} (-V_t(\Theta_t u)) = 0.$$

Hence the results follows.

A similar result holds for the Klein–Gordon $\square_g u + u^3 = 0$ on a 4-dimensional hyperbolic pseudo-Riemannian manifold, by using Theorem 3.2 and Theorem 6.1. \square

7 Comparison with quantum field theory

The space \mathbb{F} shares some analogies with the Fock spaces used by physicists in the quantum field theory. In the following we set $N := \dim E$, we let (e_1, \dots, e_N) be a basis of E and we use the affine coordinates $E \ni w \mapsto w^i \in \mathbb{R}$, for $i = 1, \dots, N$, in this basis. First assume that $s > n/2$, so that \mathcal{E}_0^s embeds continuously in continuous functions. Then for all $x \in \mathcal{M}$ and $i = 1, \dots, N$ we define the continuous linear map $\phi^i(x) : \mathcal{E}_0^s \rightarrow \mathbb{R}$ (equivalently $\phi^i(x) \in (\mathcal{E}_0^s)^* \subset \mathbb{F}$) by

$$\begin{aligned} \phi^i(x) : \mathcal{E}_0^s &\longrightarrow \mathbb{R} \\ \varphi &\longmapsto \varphi^i(x). \end{aligned}$$

If s is arbitrary we define ϕ^i as a distribution on \mathcal{M} , with values in $(\mathcal{E}_0^s)^* \subset \mathbb{F}$ by

$$\begin{aligned} \phi^i : \mathcal{C}_c^\infty(\mathcal{M}) &\longrightarrow (\mathcal{E}_0^s)^* \\ f &\longmapsto \left[\int_{\mathcal{M}} f(x) \phi^i(x) dx : \varphi \longmapsto \int_{\mathcal{M}} f(x) \varphi^i(x) dx \right] \end{aligned}$$

Similarly we define $\frac{\partial \phi^i}{\partial x^\mu}$ as a distribution with values in $(\mathcal{E}_0^s)^*$. More generally, assuming that s is s.t. we can make sense of $N(\varphi, \partial\varphi)$, we define the \mathbb{F} -valued distribution $N(\phi, \partial\phi)$. Note that the constant functional $\mathbf{1}$ equal to 1 on \mathcal{E}_0^s plays a role analogous to the vacuum.

As an algebra of functions (on \mathcal{E}_0^s) \mathbb{F} acts linearly on itself by multiplication: to each $g \in \mathbb{F}$ we associate the multiplication linear operator $[f \mapsto gf] \in \text{End}(\mathbb{F})$. This defines a natural embedding $\mathbb{F} \hookrightarrow \text{End}(\mathbb{F})$ and all previous \mathbb{F} -valued distributions $\phi^i, \frac{\partial \phi^i}{\partial x^\mu}, N^i(\phi, \partial\phi)$ can also be viewed as $\text{End}(\mathbb{F})$ -valued distributions.

Another important type of $\text{End}(\mathbb{F})$ -valued distribution is:

$$\begin{aligned} \phi_i^+ : \mathcal{C}_c^\infty(\mathcal{M}) &\longrightarrow \text{End}(\mathbb{F}) \\ f &\longmapsto \left[\int_{\mathcal{M}} f(y) \phi_i^+(y) dy : f \longmapsto \delta f_{\int_{\mathcal{M}} f(y) G_y e_i dy} \right]. \end{aligned}$$

Here, in the case where $\mathcal{M} = M$ is a flat space-time, G_y is defined by: $\forall x, y \in M, G_y(x) := G(x - y)$, where G is the distribution defined in (22). In the case where \mathcal{M} is a curved globally hyperbolic space-time and if $L = \square_g$, G_y is defined in Section 3.2, i.e. is the solution of $\square_g G_y + m^2 G_y = 0$ with the Cauchy conditions $G_y|_\sigma = 0$ and $\langle N, \nabla G_y \rangle_g|_\sigma = \delta_y$, for any Cauchy hypersurface σ which contains y . In both case it may be useful to set $G(x, y) := G_x(y)$.

Hence for any $f \in \mathcal{C}_c^\infty(M)$, $\int_M f(y) \phi_i^+(y) dy$ is the analytic first order operator associated with the constant vector field equal to $\int_M f(y) G_y e_i dy \in \mathcal{E}_0^s$ everywhere. Intuitively one may think that the notation $\phi_i^+(y)$ would represent the first order operator $f \mapsto \delta f_{G_y e_i}$ associated with the constant vector field $G_y e_i$, if $G_y e_i$ would be in \mathcal{E}_0^s (but it does not here if $s > n/2$).

This language allows us to express the operator $V_{t \cdot}$ of our Main Theorem as:

$$V_{t \cdot} := \int_{\mathbb{R}^n} d\vec{y} N^i(\phi, \partial\phi)(t, \vec{y}) \phi_i^+(t, \vec{y}) = \int_{y^0=t} d\vec{y} N^i(\phi, \partial\phi)(y) \phi_i^+(y),$$

where we assume a summation over the repeated index i . The expression (6) can be written as $U_{t_1}^{t_2}(\Theta_{t_2} u)$, where

$$U_{t_1}^{t_2} = T \exp \int_{t_1}^{t_2} dy^0 \int_{\mathbb{R}^n} d\vec{y} N^i(\phi, \partial\phi)(y) \phi_i^+(y).$$

We can then recover an expansion of this integral with terms analogous by using Wick's theorem with the commutation rules

$$[\phi^i(x), \phi^j(y)] = [\phi_i^+(x), \phi_j^+(y)] = 0, \quad [\phi_i^+(x), \phi^j(y)] = G_{ij}^j(x, y) = G_{ix}^j(y),$$

where $G_{ij}^j := (e^j, G e_i)$ (here (e^1, \dots, e^N) is the dual basis of (e_1, \dots, e_N)). In other words the $\phi_i^+(x)$'s play the role of *annihilation* operators and the $\phi^i(x)$'s play the role of *creation* operators.

As an example, we consider solutions u of the scalar equation $\square_g u + u^3 = 0$ on a 4-dimensional space-time (\mathcal{M}, g) (see Section 3) and we are given a smooth family of

admissible Cauchy hypersurfaces $(\sigma_s)_{s \in \mathbb{R}}$ which, for simplicity, we assume to be the level sets $\sigma_s = \tau^{-1}(s)$ of a temporal function $\tau \in \mathcal{C}_c^\infty(\mathcal{M})$. We let $V(s, \varphi) := \Phi_{\sigma_s}(0, \lambda_s \varphi^3|_{\sigma_s})$ be the associated family of vector fields. We can express it more intuitively by setting

$$V_s = \int_{\sigma_s} d\mu_g(y) \lambda_s(y) \phi(y)^3 G_y, \quad \text{so that} \quad V_s(\varphi) = \int_{\sigma_s} d\mu_g(y) \lambda_s(y) \varphi(y)^3 G_y.$$

Then the corresponding first order operator reads

$$V_s \cdot = \int_{\sigma_s} d\mu_g(y) \lambda_s(y) \phi(y)^3 \phi^+(y) = \int_{\sigma_s} d\bar{y} \phi(y)^3 \phi^+(y),$$

where we introduced the shorter notation $d\bar{y} := d\mu_g(y) \lambda_s(y)$. Let $f \in (\mathcal{E}_0^s)^*$ be linear, of the form $f = \int_{\mathcal{M}} d\text{vol}_g(x) \alpha(x) \phi(x)$ (or equivalently $f(\varphi) = \int_{\mathcal{M}} d\text{vol}_g(x) \alpha(x) \varphi(x)$, $\forall \varphi \in \mathcal{E}_0^s$), where $\alpha \in \mathcal{C}_c^\infty(\mathcal{M})$. Then

$$V_s \cdot f = \int_{\mathcal{M}} d\text{vol}_g(x) \alpha(x) \int_{\sigma_s} d\bar{y} G_y(x) \phi(y)^3$$

and, writing $d\text{vol}_g(x) \simeq dx$ for short,

$$V_{s_2} \cdot (V_{s_1} \cdot f) = 3 \int_{\mathcal{M}} \alpha(x) dx \int_{\sigma_{s_2}} d\bar{y}_2 \int_{\sigma_{s_1}} d\bar{y}_1 G_{y_1}(x) G_{y_2}(y_1) \phi(y_2)^3 \phi(y_1)^2.$$

We thus deduce the first terms in the expansion of $U_{t_1}^{t_2} f$ (relating the Cauchy data σ_{t_1} and σ_{t_2}).

$$\begin{aligned} U_{t_1}^{t_2} f &= f + \int_{t_1}^{t_2} ds V_s \cdot f + \int_{t_1}^{t_2} ds_2 \int_{t_1}^{s_2} ds_1 V_{s_2} \cdot (V_{s_1} \cdot f) + \dots \\ &= \int_{\mathcal{M}} \alpha(x) dx \phi(x) + \int_{t_1}^{t_2} ds \int_{\mathcal{M}} \alpha(x) dx \int_{\sigma_s} d\bar{y} G_y(x) \phi(y)^3 \\ &\quad + 3 \int_{t_1}^{t_2} ds_2 \int_{t_1}^{s_2} ds_1 \int_{\mathcal{M}} \alpha(x) dx \int_{\sigma_{s_2}} d\bar{y}_2 \int_{\sigma_{s_1}} d\bar{y}_1 G_{y_1}(x) G_{y_2}(y_1) \phi(y_2)^3 \phi(y_1)^2 + \dots \end{aligned}$$

Using $ds d\bar{y} = ds d\mu_g(y) \lambda_s(y) = d\text{vol}_g(y) \simeq dy$ and setting $\int_{\sigma_{t_1}}^{\sigma_{t_2}} dy = \int_{t_1 < \tau(y) < t_2} dy$,

$$\begin{aligned} U_{t_1}^{t_2} f &= \int_{\mathcal{M}} \alpha(x) dx \phi(x) + \int_{\mathcal{M}} \alpha(x) dx \int_{\sigma_{t_1}}^{\sigma_{t_2}} dy G_y(x) \phi(y)^3 \\ &\quad + 3 \int_{\mathcal{M}} \alpha(x) dx \int_{\sigma_{t_1}}^{\sigma_{t_2}} dy_2 \int_{\sigma_{t_1}}^{\sigma_{\tau(y_2)}} dy_1 G_{y_1}(x) G_{y_2}(y_1) \phi(y_2)^3 \phi(y_1)^2 + \dots \end{aligned}$$


Now apply Theorem 0.2: for any solution u of $\square_g u + u^3 = 0$ and for $\|u\|$ and $|t_2 - t_1|$ sufficiently small, we have $f(\Theta_{t_1} u) = (U_{t_1}^{t_2} f)(\Theta_{t_2} u)$. Hence assuming for simplicity that u is continuous, $f = \phi(x)$ for some $x \in \mathcal{M}$ and $t_1 > t_2$, we get

$$\begin{aligned} \Theta_{t_1} u(x) &= \Theta_{t_2} u(x) - \int_{\sigma_{t_2}}^{\sigma_{t_1}} dy G_y(x) (\Theta_{t_2} u(y))^3 \\ &\quad + 3 \int_{\sigma_{t_2}}^{\sigma_{t_1}} dy_2 \int_{\sigma_{\tau(y_2)}}^{\sigma_{t_1}} dy_1 G_{y_1}(x) G_{y_2}(y_1) (\Theta_{t_2} u(y_2))^3 (\Theta_{t_2} u(y_1))^2 + \dots \end{aligned} \tag{99}$$

Each term of the form $\Theta_t u(x)$ (marked by a line with a bold foot on the diagram below) reads:

$$\Theta_t u(x) = \int_{\sigma_t} d\mu_g(y) (\langle N, \nabla G_x \rangle_g(y) u(y) - G_x(y) \langle N, \nabla u \rangle_g(y)).$$

Identity (99) (for e.g. $t_2 < t_1 \leq x^0$) is pictured by the following diagram representation where Feynman rules are used (see [21]). Note that if $x^0 = t_1$, the l.h.s. of (99) is nothing



but $u(x)$.

8 A list of examples

Klein–Gordon equations

The Main Theorem can be applied to all nonlinear Klein–Gordon equations of the type $\square u + m^2 + N(u) = 0$, where u is a (real-valued) scalar field and N is a real analytic function (e.g. any polynomial or trigonometric function) for $s > n/2$. However as already stressed in Remark 2.1 this result extends straightforwardly to the case $s = 1 \leq n/2$, if N is a polynomial of degree less than or equal to $n/n - 2$.

Schrödinger equations

Our result can be applied only in the case where $n = 1$, for any real analytic nonlinear function N , i.e. to the equation $i\partial_0 u + (\partial_1)^2 u + N(u) = 0$ and for $s > 1/2$, since $H^s(\mathbb{R})$ is then an algebra and because of the continuous embedding $H^s(\mathbb{R}) \hookrightarrow L^2(\mathbb{R})$.

Wave maps

We consider for instance wave maps into the unit sphere $S^k \subset \mathbb{R}^{k+1}$ (but we may replace S^k by any Riemannian manifold which admits a real analytic isometric embedding in some Euclidean space). We set $H^s(\mathbb{R}^n, S^k) := \{v \in H^s(\mathbb{R}^n, \mathbb{R}^{k+1}); v(x) \in S^k \text{ a.e.}\}$. Wave maps are maps $u \in \mathcal{C}^0(\mathbb{R}, H^s(\mathbb{R}^n, S^k)) \cap \mathcal{C}^1(\mathbb{R}, H^{s-1}(\mathbb{R}^n, \mathbb{R}^{k+1}))$, which are weak solutions of the system:

$$\square u + (|\partial_0 u|^2 - |\vec{\partial} u|^2)u = 0.$$

We note that the nonlinearity $N(u, \partial u) = (|\partial_0 u|^2 - |\vec{\partial} u|^2)u$ is quadratic in ∂u and hence does not satisfy (20). Thus our result applies with $s > n/2 + 1$.

The Dirac–Maxwell system

Set $n = 4$, $g_{\mu\nu} := \text{diag}(1, -1, -1, -1) = g^{\mu\nu}$ and consider 4×4 Dirac matrices $\gamma^0, \gamma^1, \gamma^2, \gamma^3$ satisfying the Clifford algebra condition $\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2g^{\mu\nu}$. We agree to sum over any repeated index. The Dirac operator is $\not{\partial} = \gamma^\mu \partial_\mu$, acting on functions $\psi : \mathbb{R}^4 \rightarrow \mathbb{C}^4$. The Dirac–Maxwell system can be written

$$\begin{cases} i\not{\partial}\psi - m\psi &= e\gamma^\mu \psi A_\mu \\ \partial_\nu F^{\nu\mu} &= e\bar{\psi}\gamma^\mu \psi, \end{cases} \quad (100)$$

where $A = A_\mu dx^\mu$ is a gauge connection for the electromagnetic field, $F_{\mu\nu} := \partial_\mu A_\nu - \partial_\nu A_\mu$ is the electromagnetic field and $F^{\mu\nu} := g^{\mu\lambda} g^{\nu\sigma} F_{\lambda\sigma}$. If we further assume the Lorentz gauge condition $\partial_\mu A^\mu = 0$, where $A^\mu := g^{\mu\nu} A_\nu$, then $\partial_\nu F^{\nu\mu} = \square A^\mu$, so that (100) can be written:

$$\begin{cases} i\not{\partial}\psi - m\psi - e\gamma^\mu \psi A_\mu &= 0 \\ \square A^\mu - e\bar{\psi}\gamma^\mu \psi &= 0, \end{cases} \quad (101)$$

i.e. has the form (19). Our result can hence be applied if we assume that $A \in C^0(\mathbb{R}, H^s(\mathbb{R}^3, (\mathbb{R}^4)^*)) \cap C^1(\mathbb{R}, H^{s-1}(\mathbb{R}^3, (\mathbb{R}^4)^*))$ and $\psi \in C^0(\mathbb{R}, H^s(\mathbb{R}^3, \mathbb{C}^4)) \cap C^1(\mathbb{R}, H^{s-1}(\mathbb{R}^3, \mathbb{C}^4))$, for $s > 3/2$. Note that the Lorentz gauge can be achieved by starting from any arbitrary gauge connection \tilde{A}_μ by setting $A_\mu = \tilde{A}_\mu - \partial_\mu \varphi$, where $\varphi(x) = \int_0^{x^0} dy^0 \int_{\mathbb{R}^3} d\vec{y} G(x-y) (\partial_\mu \tilde{A}^\mu)(y)$ (so that $\square \varphi = \partial_\mu \tilde{A}^\mu$).

The pure Yang–Mills equation

Given a finite dimensional semi-simple Lie algebra \mathfrak{g} , the Yang–Mills equation for a connection $A : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathfrak{g}$ reads:

$$\partial_\nu (\partial^\nu A^\mu - \partial^\mu A^\nu + [A^\nu, A^\mu]) + [A_\nu, \partial^\nu A^\mu - \partial^\mu A^\nu + [A^\nu, A^\mu]] = 0, \quad (102)$$

where we use the same convention on repeated indices as in the previous paragraph. Assuming again that the Lorentz gauge condition $\partial_\mu A^\mu = 0$ is satisfied (which, in this nonlinear case, is harder to achieve than for electromagnetism), then the higher order term is simply $\partial_\nu (\partial^\nu A^\mu - \partial^\mu A^\nu) = \square A^\mu$. Then (102) has the form

$$\square A^\mu + \underbrace{[A_\nu, [A^\nu, A^\mu]]}_{\text{cubic in } A} + \underbrace{\partial_\nu ([A^\nu, A^\mu]) + [A_\nu, \partial^\nu A^\mu - \partial^\mu A^\nu]}_{\text{linear in } \partial A, \text{ linear in } A} = 0 \quad (103)$$

and hence satisfies Hypothesis (20). Thus Theorems 0.1 and 0.2 can be applied to solutions of (103) if $s > n/2 > s - r$, i.e., since $n = 3$ and $r = 1$, if $3/2 < s < 5/2$.

References

- [1] A. Agrachev and R. Gamkrelidze, *Chronological algebras and nonstationary vector fields*, Journal Soviet Math. (1979), 17, 1650–1675.
- [2] S Alinhac, *Hyperbolic partial differential equations*, Universitext, Springer 2009.

- [3] S. Alinhac, P. Gérard, *Pseudo-differential operators and the Nash–Moser theorem*, Graduates Studies Maths., Vol. 82, AMS.
- [4] J. Baez, *Scattering and the geometry of solution manifold of $\square f + \lambda f^3 = 0$* , J. Funct. Anal. 94 (1989), 317–332.
- [5] J. Baez, Z. Zhou, *Scattering and complete integrability in the massive φ^4 theory*, J. Funct. Anal. 83 (1990), 397–414.
- [6] C. Bär, N. Ginoux, F. Pfäffle, *Wave equations on Lorentzian manifolds and quantization*, EMS 2007.
- [7] P. Brenner, *On scattering and everywhere defined scattering operators for nonlinear Klein–Gordon equations*, J. Diff. Equa. 56 (1985), 310–344.
- [8] C. Brouder, *Runge–Kutta methods and renormalization*, Eur. Phys. J. C. 12 (2000), 521–534.
- [9] C. Brouder, *Trees, renormalization and differential equations*, B.I.T. 44 (2004), no. 6, 425–438.
- [10] J.C. Butcher, *The numerical analysis of ordinary differential equations*, Wiley, Chichester, 1987.
- [11] R. Carles, I. Gallagher, *Analicity of the scattering operator for semilinear dispersive equations*, Commun. Math. Phys. 286 (2009), 1181–1209.
- [12] K.T. Chen, *Integration of paths, geometric invariants and a generalized Baker–Hausdorff formula*, Annals of Math. 65 (1957), 163–178.
- [13] Y. Choquet-Bruhat, D. Chistodoulou and M. Francaviglia, *On the wave equation on curved space-time*, Ann. IHP, Sec. A, T. 31, n. 4 (1979), 399–414.
- [14] M. Dütsch, K. Fredenhagen, *The master Ward identity and generalized Schwinger–Dyson equation in classical field theory*, Commun. Math. Phys. 243 (2003), no.2, 275–314.
- [15] F. Finster, J. Tolksdorf, *Bosonic Loop Diagrams as Perturbative Solutions of the Classical Field Equations in φ^4 -Theory*, J. Mathematical Physics 53 (2012).
- [16] M. Fliess, *Fonctionnelles causales non linéaires et indéterminées non commutatives*, Bull. Soc. Math. France 109 (1981), no. 1, 3–40.
- [17] F. Friedlander, *The wave equation on a curved space-time*, Cambridge University Press, Cambridge, 1975.
- [18] E. Hairer and G. Wanner, *On the Butcher group and general multi–value methods*, Computing (1974), 13 (1), 1–15.

- [19] D. Harrivel, *Planar binary trees and perturbative calculus of observables in classical field theory*, Ann. IHP 23 (2006), 891–909.
- [20] D. Harrivel, *Butcher series and control theory*, preprint arXiv math/0603133
- [21] D. Harrivel, F. Hélein, *First integrals for nonlinear hyperbolic equations*, arXiv:0704.2674
- [22] E. Hebey, *Sobolev spaces on Riemannian manifolds*, Springer LMS 1635, Springer 1996.
- [23] F. Hélein, *Harmonic maps, conservation laws and moving frames*, Cambridge tracts in Mathematics 150, CUP 2002.
- [24] F. Hélein, *Hamiltonian formalisms for multidimensional calculus of variations and perturbation theory*, in *Noncompact problems at the intersection of geometry, analysis, and topology*, Contemp. Math., 350 (2004), 127–147.
- [25] F. Hélein, *Multisymplectic formalism and the covariant phase space*, in *Variational Problems in Differential Geometry*, K. Houston, R. Bielawski, M. Speight, Eds., London Mathematical Society Lecture Note Series 394, Cambridge University Press, 2012, p. 94–126.
- [26] M. Kawski, H. Sussman, *Noncommutative power series and formal Lie-algebraic techniques in nonlinear control theory*, in *Operators, Systems and linear algebra*, 1997, 111–128.
- [27] D. Kreimer, *On the Hopf algebra structure of perturbative quantum field theory*, Adv. Th. Math. Phys., 2 (1998), pp. 303–334.
- [28] J. Leray, *Hyperbolic differential equations*, Princeton, I.A.S. ed., 1952.
- [29] W. Magnus, *On the exponential solution of differential equations for a linear operator*, Comm. Pure Appl. Math. (1954), VII, 649–673.
- [30] C. Morawetz, W. Strauss, *On a nonlinear scattering operator*, Comm. Pure Appl. Math. 26 (1973), 47–54.
- [31] P. Racka, W. Strauss, *Analyticity properties of the scattering operator in nonlinear relativistic classical and prequantized field theories*, Rep. Math. Phys. 16 (1979), 317–327.
- [32] H. Sussmann, *A general theorem on local controllability*, SIAM J. Control Optim. 25 (1987), no. 1, 158–194.
- [33] W. Strauss, *Everywhere defined wave operators*, in “Nonlinear Evolution Equations” (M. Crandall, Ed.), p. 85–101, Academic Press, New York, 1978.
- [34] W. Strauss, *Nonlinear scattering at low energy*, J. Funct. Anal. 41 (1981), 110–133.